


- 30.** Sollner L, Shaqireen DOK, Wu JT and Schwarz H. Signal transduction mechanisms of CD137 ligand in human monocytes. *Cell Signal*. 2007; 19:1899-1908.
- 31.** Kwon BS, Hurtado JC, Lee ZH, Kwack KB, Seo SK, Choi BK, Koller BH, Wolisi G, Broxmeyer HE and Vinay DS. Immune responses in 4-1BB (CD137)-deficient mice. *J Immunol*. 2002; 168:5483-5490.
- 32.** Pollok KE, Kim YJ, Hurtado J, Zhou Z, Kim KK and Kwon BS. 4-1BB T-cell antigen binds to mature B cells and macrophages, and costimulates anti-mu-primed splenic B cells. *Eur J Immunol*. 1994; 24:367-374.
- 33.** Pauly S, Broll K, Wittmann M, Giegerich G and Schwarz H. CD137 is expressed by follicular dendritic cells and costimulates B lymphocyte activation in germinal centers. *J Leukoc Biol*. 2002; 72:35-42.
- 34.** Middendorp S, Xiao Y, Song JY, Peperzak V, Krijger PH, Jacobs H and Borst J. Mice deficient for CD137 ligand are predisposed to develop germinal center-derived B-cell lymphoma. *Blood*. 2009; 114:2280-2289.

## SUPPLEMENTARY FIGURE



Supplementary Figure 1.