

Origin and pathophysiology of protein carbonylation, nitration and chlorination in age-related brain diseases and aging

Efstathios S. Gonos¹, Marianna Kapetanou^{1,2}, Jolanta Sereikaite³, Grzegorz Bartosz⁴, Katarzyna Naparto⁵, Michalina Grzesik⁵, Izabela Sadowska-Bartosz⁵

¹National Hellenic Research Foundation, Institute of Biology, Medicinal Chemistry and Biotechnology, Athens 11635, Greece

²Department of Biochemistry and Molecular Biology, Faculty of Biology, University of Athens 15701 Athens, Greece

³Department of Chemistry and Bioengineering, Faculty of Fundamental Sciences, Vilnius Gediminas Technical University, 2040 Vilnius, Lithuania

⁴Department of Molecular Biophysics, Faculty of Biology and Environmental Protection, University of Lodz, 90-236 Lodz, Poland

⁵Department of Analytical Biochemistry, Faculty of Biology and Agriculture, University of Rzeszow, 35-601 Rzeszow, Poland

Correspondence to: Izabela Sadowska-Bartosz; email: isadowska@poczta.fm

Keywords: oxidative stress, carbonylation, nitration, chlorination, proteasome

Received: April 9, 2018 **Accepted:** May 8, 2018 **Published:** May 17, 2018

Copyright: Gonos et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Non-enzymatic protein modifications occur inevitably in all living systems. Products of such modifications accumulate during aging of cells and organisms and may contribute to their age-related functional deterioration. This review presents the formation of irreversible protein modifications such as carbonylation, nitration and chlorination, modifications by 4-hydroxynonenal, removal of modified proteins and accumulation of these protein modifications during aging of humans and model organisms, and their enhanced accumulation in age-related brain diseases.

INTRODUCTION

Aging, an inevitable part of the life process, is characterized by a progressive decline in physiological functions that ultimately leads to morbidity and mortality. Aging increases susceptibility to certain class of diseases. Age-related diseases constitute a considerable socio-economic burden for contemporary societies. As human mean lifespan increases, growing incidence of these diseases has features of a pandemic. The number of people aged 65 or older is projected to grow from an estimated 524 million in 2010 to almost 1.5 billion in 2050, mostly in underdeveloped and developing countries [1]. These trends have obvious serious social and economic implications, such as healthcare costs [2].

Despite extensive studies, the molecular basis of physiological aging is still poorly understood. Reactive oxygen species (ROS), reactive nitrogen species (RNS) as well as reactive halogen species (RXS) species are believed to play a key role in the aging process. They are generated during aerobic metabolism in living organisms. The term “*reactive oxygen species*” includes both free radicals [molecules having an odd electron, like superoxide radical anion ($O_2^{\cdot-}$) and hydroxyl radical (HO^{\cdot})] and species that are not free radicals, like hydrogen peroxide (H_2O_2), singlet oxygen (1O_2) and ozone (O_3). The primary source of RNS is usually the nitric oxide radical ($\cdot NO$). In consequence of ROS and RNS reactions, peroxyxynitrite $ONOO^{\cdot}$, anion of peroxyxynitrous acid $ONOOH$, may be formed via the near

diffusion-limited reaction of $\cdot\text{NO}$ and $\text{O}_2\cdot^-$. The term “*reactive nitrogen species*” includes also nitrous acid (HNO_2), dinitrogen trioxide (N_2O_3), nitrosyl anion (NO^-), nitrosyl cation (NO^+), nitrogen dioxide radical (NO_2), peroxyxynitrate (ONOOO^\cdot), peroxyxynitric acid (ONOOOH), nitryl chloride (NO_2Cl), and nitronium cation (NO_2^+) [3, 4]. “*Reactive halogen species*” include HOCl , HOBr , HOI , chlorine, bromine, iodine etc. Hypohalogenous acids (HOX ; $\text{X} = \text{F}, \text{Cl}, \text{Br}, \text{or I}$) are formed in the body mainly by oxidation of halogen ions by myeloperoxidase. The imbalance between ROS, RNS and RXS production and the antioxidant defense, in favor of prooxidants, is causes oxidative, nitr(os)ative and halogenative stress (OS, NS, XS), respectively. Although at physiological concentrations ROS, RNS and RXS can function as signaling molecules regulating cell proliferation, growth, differentiation and apoptosis [5, 6] they react with and damage all classes of endogenous macromolecules including proteins, nucleic acids, lipids and carbohydrates [7]. Proteins are the main targets for such modifications as they are the most abundant cell components in the terms of mass content. The level of protein damage increases under stress conditions and can be in principle an integrative measure of the exposure to OS, NS and XS. However, protein turnover complicates this issue, the more that modified proteins in most cases are subject to preferential degradation [8]; see Chapter “Removal of modified proteins”.

Protein modifications produced by ROS, RNS and RXS can be classified as transient, reversible or irreversible. Reactions of free radicals with proteins leads to formation of protein radicals, which are generally short-lived, transient and are not useful as biomarkers. Protein hydroperoxides formed upon reactions with ROS are also unstable and decompose forming more stable products [9, 10]. Examples of reversible modifications are cysteine (Cys) thiol oxidation to sulfenic acid, methionine (Met) oxidation to methionine sulfoxide or cysteine *S*-nitrosylation and *S*-glutathionylation (Table 1, Fig. 1). While these modifications are of vital importance for regulation of protein function and metabolic processes, they are of less importance as permanent markers of OS/NS/XS, so this review will concentrate on irreversible protein modifications.

FORMATION OF NON-ENZYMATICALLY MODIFIED PROTEINS

Compared to other oxidative modifications, carbonyls are relatively difficult to induce and in contrast to, for example, methionine sulfoxide and cysteine disulfide bond formation, carbonylation is an irreversible oxidative process [11]. Protein carbonylation is an

oxidative modification induced by ROS, RNS, RXS and reactive aldehydes. It consists in formation of reactive aldehyde or ketone residues on proteins, which can react with 2,4-dinitrophenylhydrazine (DNPH) forming hydrazones. There are two ways of protein carbonylation. “Primary protein carbonylation” is due to oxidation of some amino acid residues, initiated by ROS, RNS and RXS, often catalyzed by metals while “secondary protein carbonylation” is caused by addition of aldehydes. The aldehydes are formed mainly in the process of lipid peroxidation [malondialdehyde, MDA; 4-hydroxy-2,3-*trans*-nonenal, (4-HNE); 2-propenal (acrolein, ACR)], but may be also by-products of glycolysis and the glycation process (methylglyoxal, glyoxal).

In the first pathway, ROS, RNS and RXS directly attack the protein producing, eventually, highly reactive carbonyl derivatives by oxidation of the side chains of lysine (Lys), arginine (Arg), proline (Pro), and threonine (Thr) residues, particularly *via* metal-catalysed oxidation, from the cleavage of peptide bonds in the α -amidation pathway or by oxidation of glutamyl residues. The main carbonyl products of metal-catalysed protein oxidation are glutamic semialdehyde, a product of oxidation of Arg, amino adipic semialdehyde, a product of Lys oxidation, 2-pyrrolidine, a product of histidine (His) oxidation and 2-amino-3-ketobutyric acid, a product of oxidation of Thr (Fig. 1E) [12]. Carbonylation is site-specific; an iterative statistical method has been proposed to identify potential sites of carbonylation [13].

The second type of reaction involves the addition of reactive aldehyde groups to the side chains of Cys, His, or Lys residues *via* Michael addition (Fig. 1F). Reactive carbonyl groups can be also generated through the reaction of the amino group of Lys residues with reducing sugars or their oxidation products (glycation/glycoxidation products) [14].

Dimerization of tyrosyl radicals (Tyr^\cdot) leads to the formation of dityrosine (Fig. 1H). Products of oxidative destruction of tryptophan (Try) include kynurenine and N-formylkynurenine (Fig. 1I). All these products have their characteristic fluorescence and their content can be easily evaluated fluorimetrically [15, 16].

RNS can oxidize proteins and alter their biological functions also in other ways. Nitration of amino acids, such as tyrosine (Tyr) and, to a lesser extent, Try and His, is an important form of protein modification that occurs during NS [17]. Tyr, a nonessential aromatic amino acid, carrying a hydroxyl group, is often exposed hem vulnerable to nitration, as well as oxidation [18, 19].




Figure 1. Selected non-enzymatic protein modifications. (A) oxidation of cysteine residues in proteins. Cysteine residues may be oxidized to sulfenic, sulfinic and sulfonic derivatives or form disulfide bonds. Oxidation to sulfenic acid and formation of disulfides is reversible; (B) modifications of cysteine residues in proteins: formation of nitrosocysteine and S-glutathionylation; (C) oxidation of methionine forms methionine sulfoxide, which may be reduced back to methionine by methionine sulfoxide reductases (MSR); (D) formation of hydroperoxides of valine, lysine and leucine; (E) formation of carbonyl derivatives of lysine, arginine, His and threonine; (F) formation of 4-hydroxynonenal adducts of cysteine, His and lysine; (G) oxidative modifications of phenylalanine; (H) modifications of tyrosine; (I) modifications of tryptophan.

The nitration of Tyr is mediated by RNS such as ONOO⁻/ONOOH and [•]NO₂ although nitration can also be accomplished by heme peroxidases and nitrite [20]. The two main mechanisms of biological nitration, the ONOO⁻/ONOOH and the heme peroxidase pathways, lead both to the formation of Tyr[•] and [•]NO₂, which combine with diffusion controlled rates to form 3-nitrotyrosine (3-NT; Fig. 1H). The oxidants leading to Tyr[•] formation include CO₃^{•-}, [•]OH or oxo-metal complexes. Importantly, [•]NO₂ alone is inefficient in promoting nitration, because its reaction with Tyr to produce Tyr[•] is slow compared to other processes that [•]NO₂ undergoes. I. a., reaction with another Tyr[•] to form

3,3-dityrosine competes with the formation of 3-NT. However, under certain conditions protein radicals can be stabilized, e. g. when intra- and intermolecular dimerization is limited due to diffusional and spatial constraints, both in aqueous and hydrophobic compartments. In such cases reaction of Tyr[•] with [•]NO₂ may be favoured. Another pathway competing with Tyr nitration is the formation of 3-hydroxytyrosine, which can be performed mainly by [•]OH or oxo-metal complexes. An alternative radical mechanism for Tyr nitration involves the reaction of a Tyr[•] with [•]NO to form 3-nitrosotyrosine followed by two-electron oxidation to 3-NT [21].

Table 1. Most important oxidative, nitrative and chlorinative modifications of proteins. After [11], modified.

Amino acid	Modification	Stability/Reversibility
Cysteine	Oxidation of –SH to sulfenic acid (-SOH), sulfinic acid (-SO ₂ H) or sulfonic acid (-SO ₃ H) Formation of a disulfide bond –SS–	First stage, and in some cases second stage reversible Reversible
Cysteine	Nitrosylation [formation of (-SNO)]	Reversible
Cysteine	Glutathionylation	Reversible
Tyrosine, tryptophan, other amino acids	Protein radicals	May be reduced or react to form further products
Glutamic acid, tyrosine, lysine, leucine, valine, proline, isoleucine	Hydroperoxides	May be reduced; decompose to further products
Histidine	2-Oxohistidine	Irreversible
Lysine, arginine, proline, threonine	Formation of carbonyl derivatives by direct oxidative attack on amino-acid side chains (α -amino adipic semialdehyde from lysine, glutamic semialdehyde from arginine, 2-pyrrolidone from proline, and 2-amino-3-ketobutyric acid from threonine)	Irreversible [Decarbonylation?]
Lysine, cysteine, histidine	Formation of carbonyl derivatives by secondary reaction with reactive carbonyl compounds derived from oxidation of carbohydrates (glycoxidation products), lipids (MDA, 4-HNE, ACR) and advanced glycoxidation and lipoxidation end products	Irreversible
Methionine	Methionine sulfoxide	Reversible by methionine sulfoxide reductases
Phenylalanine	<i>o</i> -Tyrosine, <i>m</i> -tyrosine	Irreversible
Tyrosine	Hydroxylation to 3,4-dihydroxyphenylalanine Dimerization to dityrosine	Irreversible
Tyrosine, tryptophan, histidine	Nitration [introduction of (-NO ₂)]	Irreversible [Denitration ?]
Tyrosine	Chlorination to 3-chlorotyrosine	Irreversible
Tryptophan	5-Hydroxytryptophan, 7-hydroxytryptophan, kynurenine, N-formylkynurenine	Irreversible

Hypochlorous acid (HOCl) is the main player involved in protein chlorination *in vivo* [16]. HOCl is generated by the reaction of H₂O₂ with chloride ions (Cl⁻) catalysed by myeloperoxidase (MPO, EC 1.11.1.7) [22-24]. For a long time, myeloperoxidase (MPO) was regarded as the only human enzyme known to produce HOCl at the physiological concentrations of chloride (100-140 mM) [25]. Nevertheless, recent findings revealed that another mammalian heme peroxidase, peroxidase 1, is capable of catalysing the oxidation of chloride to HOCl, too. The enzyme is also known as vascular peroxidase 1 [26-29]. Up to 80% of the H₂O₂ generated by activated neutrophils may be used to produce local concentrations as high as 20-400 μ M

HOCl within an hour [30, 31]. The pK_a of HOCl is 7.59 [32], so at physiological pH values, HOCl exists in equilibrium with its anion ⁻OCl at approximately equal concentrations. HOCl is a powerful oxidant and plays an important physiological role. MPO-produced HOCl is involved in innate immune response and kills invading pathogens [33, 34]. Green et al. [35] showed that the diminution of HOCl production observed with decreasing Cl⁻ availability results in impaired killing of bacteria. However, during chronic inflammation the excessive production of HOCl leads to the host tissue damage and plays a pathophysiological role in inflammatory diseases [36]. Proteins are major targets for HOCl, and the reactions of this oxidant with proteins

result in side-chain modifications (mainly chlorination of Tyr residues, Fig. 1H), cross-linking and backbone fragmentation [37, 38].

PROTEIN CARBOXYLATION IN AGING AND AGE-RELATED DISEASES

Protein carbonyl content is the most general and broadly used biomarker of oxidative protein damage and, more generally, OS. However, protein carbonyls are important not only as a biomarker for protein oxidation in aging and disease. They have also been shown to impair protein structure and function and to participate in the etiology and progress of diseases and age-related changes in the body [39, 40]. Carbonylation may alter the conformation of the polypeptide chain, which leads to partial or total inactivation of proteins. The consequent loss of function or structural integrity of carbonylated proteins can have a wide range of downstream functional consequences and may underlie the subsequent cellular dysfunctions and tissue damage [41]. Protein carbonylation was demonstrated to modify activities of enzymes and other protein functions like DNA binding of transcription factors [42]. Carbonylation can lead to functional impairment of proteins involved in insulin signaling, so the insulin signaling pathway gets disrupted by carbonylation [43].

Another mechanism of protein carbonyl action involves inhibition of proteasomal activity. While moderately carbonylated proteins are degraded by the proteasomal system, heavily carbonylated proteins form high-molecular-weight aggregates that are not digested and accumulate. Such aggregates of carbonylated proteins are resistant to degradation and can inhibit proteasomes. Neurodegenerative diseases are directly associated with the accumulation of proteolysis-resistant aggregates of carbonylated proteins in tissues [39].

It has been hypothesized that protein carbonylation is reversible, and protein carbonyls can be removed by a "decarbonylase" activity; thus, protein carbonylation can play a role in cellular signaling. Thioredoxin was postulated to be involved in protein decarbonylation [44-46]. However, experimental support for this hypothesis is scarce. Age-related increase in the protein carbonyl content has been demonstrated in many objects. Data from various laboratories demonstrated a dramatic increase in the content of carbonylated proteins during the last third of the lifespan of various objects, i. a., human dermal fibroblasts in culture [47], human lens [48], rat liver [49], house fly [50] and *Caenorhabditis elegans* [51]. Further examples of age-related increase in the level of protein carbonyls are given in Table 2.

Table 2. Examples of studies on the effect of aging on protein carbonyl level.

Problem studied	Material or object studied/methods	Findings	Reference
Effect of replicative aging of fibroblasts <i>in vitro</i> on protein carbonyl level	WI-38 fibroblasts, intermediate or middle-aged (PD between 25 and 39) and replicatively senescent (PD < 40)/OxyBlot	Increase in the level of carbonylated proteins, Preferential carbonylation of certain proteins	[52]
Effect of replicative aging and heat stress on protein carbonyl level in human fibroblasts	Human foreskin fibroblasts, middle-aged and senescent/OxyBlot	Increased protein carbonyl content in senescent cells and in heat stressed cells, without recovery	[53]
Effect of aging and late onset dietary restriction on the protein carbonyl level in cerebral hemispheres	BALB/c mice, 4-w old and 84-w old/DNPH assay and WB	Increased protein carbonyl level in old mice, Reduction of protein carbonyl level after 3-m calorie restriction	[54]
Effect of aging on protein carbonyl level of high-molecular weight protein aggregates isolated from the bone marrow and 5	Female C57BL/6 J mice/Oxyblot, Protein Carbonyl Assay kit	Enhanced protein carbonyl level in 22-m old vs 3-m and 12-m old mice	[55]
Effect of aging on protein carbonyl level of testis mitochondria	5-m vs 30-m old rats/2D PAGE, WB, carbonyl detection with biotin-hydrazide	Age-related increase in the carbonyl content of many proteins, decrease for some proteins	[56]

Comparison of protein carbonyl content in mitochondria of slow-twitch and fast-twitch muscles	Fisher 344 female rats/MS	Fast-twitch muscle contain twice as many carbonylated mitochondrial proteins as slow-twitch muscle	[57]
Effect of age on protein carbonyl content of erythrocyte membranes	49 healthy subjects of both sexes aged 17- 80 y/DNPH assay	High correlation between age and protein carbonyl content, Negative correlation between protein carbonyl content and total antioxidant capacity of plasma (FRAP)	[58]
Effect of age on carbonyl content of mouse liver	Young (3 m) and aged (24 m) male C57BL/6 mice/2D PAGE, WB	Increased protein carbonylation in aged mice, especially of BiP/Grp78, protein disulfide isomerase (PDI) and calreticulin	[59]
Effect of age on protein carbonyl level in rat cerebral cortex and hippocampus	4-m, 12-m and 22-m old rats/DNPH assay	Higher protein carbonylation in hippocampus than in cerebral cortex, Increase in protein carbonyl level with age, attenuated by physical exercise	[60]
Effect of age and sarcopenia on carbonyl content of skeletal muscle subfractions	Sarcoplasmic, myofibrillar, and mitochondrial subfractions from musculus vastus lateralis biopsies of 16 young and 16 elderly persons/Oxyblot	Increased mitochondrial (but not myofibrillar or sarcoplasmic) protein carbonyl content with aging, No effect of stage I sarcopenia	[61]
Effect of age and gender on protein carbonyl content in saliva and plasma	273 healthy Chinese subjects, aged between 20 and 79/ELISA	Significant correlation of saliva and plasma protein carbonyls with age, No relation to gender	[62]
Effect of aging on protein carbonyl level of mouse skeletal muscles	Muscles from 3, 15, 24, 27 and 29 m old female C57Bl/6J mice/DNPH assay	Protein carbonyl level of gastrocnemius muscles unchanged between 3 and 15 m, increasing at 27 and 29 m. No age-related decrease in protein thiol level or increase in the levels of MDA and F2-isoprostanes	[63]
Effect of age and physical exercise on the carbonyl level of plasma proteins	481 participants of both sexes aged 65-69 y and 239 participants aged 90 y or more/ELISA	Elevation of protein carbonylation with aging, attenuated by physical activity	[64]
Effect of aging on the level of protein carbonyls in human rectus abdominis and vastus lateralis muscles	Muscle biopsies of 11 children 0-12 y old and 11 persons 52-76 y old/2D PAGE, WB	No significant differences in the global level of protein carbonyls between the groups in both rectus abdominis and vastus lateralis muscles	[65]
Effect of age on protein carbonyl content of external intercostals and quadriceps muscles	12 young and 12 elderly persons of both sexes/ DNPH assay	Increased levels of protein carbonyls in external intercostals of elderly women, but not of elderly men	[66]

DNPH, dinitrophenylhydrazine; 2D PAGE, two-dimensional polyacrylamide gel electrophoresis; MS, mass spectrometry, PD, population doublings; WB, Western Blotting

Many studies have demonstrated the existence of a relationship between the level of protein carbonylation in cells or tissues/organs and human or animal age and lifespan, as exemplified below. The level of protein carbonylation in human fibroblasts have been found to increase exponentially with advancing age of donors [67]. When mobility of fruit flies *Drosophila melanogaster* was restricted by culturing under conditions preventing flying, the lifespan of the insects was increased 2 to 3-fold. Age-related accumulation of protein carbonyls was slower in mobility-restricted flies in comparison with control flies, which could fly and had shorter lifespan [50]. Transgenic fruit flies which overexpressed CuZn-superoxide dismutase (SOD1) and catalase had prolonged lifespan. Accumulation rate of protein carbonyls in these flies was slower than in control flies whose lifespan was shorter [68]. Calorie restriction was also demonstrated to slow down the rate of protein carbonyl accrual. In calorie-restricted mice which had lifespan increased by 35%, the rate of accumulation of protein carbonyls in several tissues was decelerated [69, 70].

Protein carbonylation is selective with respect to the protein and this rule refers also to carbonylation during aging and in age-related diseases. Approximately 10% of the proteome is more prone to carbonylation during ageing or disease than other proteins [71, 72]. In the brains of patients affected with Alzheimer's disease (AD) and Parkinson's disease (PD), mitochondrial MnSOD superoxide dismutase (SOD2) is one of the major targets of oxidative damage [73]. In turn, aconitase was found to be the only protein in the mitochondrial matrix that exhibited an age-associated increase in carbonylation in *D. melanogaster*. The accumulation of carbonyl groups was accompanied by an approximately 50% loss in aconitase activity [74]. Interestingly, the set of proteins that become carbonylated differs in various species. For example, aging-associated protein carbonylation was only seen in two proteins in mouse blood plasma, albumin and transferrin, while in the rat plasma, only albumin and α -macroglobulin showed significant progressive age-dependent carbonylation [75]. There are several possible explanations for this specificity of carbonylation. One is the presence of a transition metal on the protein, another being the localisation of proteins to be carbonylated close to ROS generating sites. However, in general the molecular basis for the apparent specificity of protein carbonylation still remains unclear [76].

Elevation of the protein carbonyl level has been reported for many diseases. Protein carbonylated levels are widely used index to determine the extent of oxidative modification of proteins both under *in vivo*

and *in vitro* conditions. Increased protein carbonyl levels were found in the cerebrospinal fluid of patients with multiple sclerosis [77] and in blood plasma of patients with multiple sclerosis [16] as well as myasthenia gravis [78]. In multiple sclerosis, increased level of protein carbonyls was also found in the brain white and gray matter [79].

The large volume of literature and heterogeneity of results makes a comprehensive understanding of the changes occurring in human brain in AD elusive. In AD, the increase in protein carbonylation level was different in various regions of the brain. It was increased by 42% in hippocampus and by 37% in the inferior parietal lobule, with respect to cerebellum, which shows little degenerative changes in this disease [80]. Some specifically carbonylated proteins in AD brain were identified in different stages of the process, including the exacerbate mild cognitive impairment (MCI) and early AD stages [81-84].

What's more, a new meta-analysis defines the pattern of changes in OS related markers by brain region in human AD and MCI brain tissue. Protein carbonylation was significantly increased in the occiput and in the hippocampus in AD, while there were no significant changes noted in other brain regions [85]. Shen et al. evaluated the levels of total protein carbonyls and identified the oxidative modification proteins in the sera of 3 \times transgenic AD mice. Their results suggested that OS is an early event in the development of AD, and analysis of specific serum protein oxidation may be more plausible for the search of AD biomarkers [86]. Brain samples of patients with Huntington disease showed increased carbonylation of more than a dozen of proteins including glial fibrillary acidic protein, aconitase, enolase 1 and creatine kinase B, glycolytic enzymes and mitochondrial proteins related to ATP production [87]. Increased level of protein carbonyls was also found in the spinal cord of G93A-SOD1 transgenic mice, an animal model of amyotrophic lateral sclerosis (ALS) [88].

Numerous data point to the role of RCS as both propagators and products of oxidative damage in neurodegenerative diseases, especially in AD [89]. In AD, the concentration of free 4-HNE was reported to be increased in the plasma and cerebrospinal fluid [90], whereas ACR content was higher in the amygdala and hippocampus/parahippocampal gyrus of AD patients [91]. MDA accumulation has been detected in the cytoplasm of astrocytes and neurons in both normal ageing and in AD patients [92]. Increased concentrations of MDA (in blood plasma and serum) and 4-HNE (in the plasma and cerebrospinal fluid) have also been reported for PD patients [91]. 4-HNE levels are

significantly elevated in the sera and spinal fluid of ALS patients and positively correlate with the extent of the disease but not with the rate of progression, which suggest 4-HNE and carbonyl bearing 4-HNE-protein adducts as possible biomarkers of the disease [77, 93].

Methods of protein carbonylation analysis

The generally used method of quantifying carbonyl groups is based on the reaction with 2,4-dinitrophenylhydrazine (DNPH). This compound reacts with carbonyl groups, forming the stable 2,4-dinitrophenylhydrazone. Dinitrophenyl group (DNP) adduct can be detected by different methods. The DNP group itself absorbs ultraviolet light, so the total carbonyl content of a protein or mixture of proteins can be quantified by a spectrophotometric assay [94]. Alkalinization of the medium may bring absorption maximum of the DNP group from 370 nm (UV) to 450 nm (visible region) [95].

The assay of carbonylated proteins has been simplified by the availability of commercial antibodies specific for DNP, which allow for their detection by immunoblotting. Dot blot analysis allows for a very sensitive quantification of the total level of protein carbonylation in a sample [96]. Immunoblotting assays based on the use of anti-DNP antibodies have been developed as an attempt to identify oxidatively damaged proteins in human tissues and body fluids. The carbonyl content in individual proteins is estimated by one-dimensional (1D) or two-dimensional (2D) sodium dodecyl sulfate (SDS) gel electrophoresis followed by Western blot immunoassay (Oxyblot). These two methods have significantly higher sensitivity and specificity than all other total carbonyl assays, but are still semiquantitative [11]. An alternative to immunochemical detection of DNP derivatives of carbonylated proteins is the reaction with a fluorescent reagents reacting with carbonyl groups such as fluorescein-5-thiosemicarbazide [97] or fluorescent hydrazides [98, 99]. Fluorescent hydrazides such as coumarin hydrazine were also used for detection of protein carbonyls in living cells [100].

DNP assay of protein carbonyls can be also combined with protein fractionation by high-performance liquid chromatography (HPLC) to obtain better sensitivity and specificity than measuring total carbonyls in a protein mixture [11]. Mass spectrometry (MS) allows for precise identification of carbonylated proteins and characterization of the carbonylation sites [101]. Proteomic tools provide a promising way to decode disease mechanisms at the protein level and help to understand how carbonylation affects protein structure and function. Recently, Havelund et al. (2017) proposed a peptide-centric approach for identification and

characterization of up to 14 different types of carbonylated amino acids in proteins. The use of diagnostic biotin fragment allows MS/MS data analysis to pinpoint sites of biotin labeling and improve the confidence of carbonyl peptide assignments [102].

NITRATIVE PROTEIN MODIFICATIONS IN AGING AND AGE-RELATED DISEASES

It should be noted that protein Tyr nitration is observed *in vivo* in healthy tissues, indicating that there is a basal flux of RNS; nevertheless, physiological nitration levels are typically low. Possible biochemical consequences of protein Tyr nitration involve changes in activity (usually loss, but sometimes gain of function), induction of immune responses, interference with tyrosine-kinase-dependent pathways, alteration of protein assembly and polymerization, and effects of protein turnover: either facilitation of protein degradation or induction of formation of proteasome-resistant protein aggregates, depending on the dose [103, 104].

Furthermore, protein Tyr nitration is also associated with physiological aging and pathophysiology of several age-related diseases such as atherosclerosis, multiple sclerosis, AD, PD, ALS, cystic fibrosis, asthma, lung diseases, myocardial malfunction, stroke, chronic hepatitis, cirrhosis, diabetes, etc. [105]. Increased content of nitrates, nitrites, and free 3-NT were also found in the cerebrospinal fluid of subjects with neurodegenerative diseases and have been proposed as functional biomarkers of neurodegeneration [16, 106].

Two faces of ^{*}NO: implications for brain aging

Nitric oxide and other RNS appear to play crucial roles in the brain such as neuromodulation, neurotransmission and synaptic plasticity, but are also involved in pathological processes such as neurodegeneration and neuroinflammation. Nitric oxide is a short-lived gaseous physiological messenger, which is highly diffusible and lipophilic in nature [107]. As an important neurotransmitter and signaling molecule, ^{*}NO is involved in numerous physiological processes throughout the nervous system. Apart from guanylate cyclase, ^{*}NO targets include ion channels, which are involved in setting neuronal excitability and calcium homeostasis, additional to its involvement in physiological plasticity processes (long-term potential-tion; long-term depression), which can include the N-methyl-D-aspartate receptor-mediated calcium-dependent activation of neuronal ^{*}NO synthase [108].

Nevertheless, ^{*}NO possesses a controversial effect on cell viability by acting both in protection against

apoptogenic stimuli, and by inducing apoptosis when produced at elevated concentrations. Moreover, excessive generation of NO , favors the formation of reactive ONOO \cdot /ONOOH and NO $_2$ species that can mediate nitration of aging brain proteins [109]. Acute and chronic inflammation result in increased NO formation and NS. It is well documented that NO and its toxic metabolite, ONOO \cdot /ONOOH, can inhibit components of the mitochondrial respiratory chain leading to cellular energy deficiency and, eventually, to cell death. Within the brain, the susceptibility of different brain cell types to NO and ONOO \cdot /ONOOH exposure may be dependent on factors such as the intracellular glutathione (GSH) concentration and cellular stress resistance signal pathways [110].

NO toxicity: regulation by glutathione

GSH is the most abundant low molecular weight thiol in mammalian cells and acts as the major cellular antioxidant. Aquilano et al. [111] reported that GSH may constitute the most important buffer of NO toxicity in neuronal cells, and demonstrated that the disruption of cellular redox buffering controlled by GSH makes neuronal cells susceptible to endogenous physiological flux of NO . GSH levels in the brain decline progressively during aging and in neurodegenerative disorders, such as AD or PD [112, 113]. It has been proposed that the decrease in GSH concentration could be mainly a consequence of the formation of protein mixed disulfides. The intracellular depletion of GSH can induce cellular stress in NO -producing cells through a NO -dependent mechanism, resulting in such effects as induction of DNA damage, inhibition of cytochrome *c* oxidase activity, accumulation of S-nitrosocysteine and increased nitration of protein Tyr residues. What's more, NO seems to be the main mediator of cell proliferation arrest through the extracellular signal-regulated kinase-1/2-p53 signaling pathway and apoptosis through the translocation of mitochondrial apoptosis-inducing nuclear factor [111].

Tyrosine nitration of synaptic proteins

Synaptic proteins can undergo extensive posttranslational modifications. Numerous evidence suggests that aging and diseases can induce nitrative stress via excessive NO production. NS can lead to uncontrolled S-nitrosylation/Tyr nitration, which can represent crucial pathological features that contribute to the onset and progression of various neurodegenerative diseases, including AD or PD [110, 111].

It has been suggested that phosphorylation and nitration of protein Tyr residues plays a role in signaling path-

ways at the nerve terminal and affects functional properties of proteins involved in the synaptic vesicle (SV) exo-endocytotic cycle [114]. Protein conformational changes induced by NO have strong impacts on protein-protein interactions in the docking/fusion steps of vesicle release. Depending on the concentration of NO and the reversibility of protein nitration, the consequences for neuronal signaling are important and relevant in physiology and pathology. According to Di Stasi and coworkers [115], ONOO \cdot /ONOOH causes Tyr nitration of SNAP-25 and Munc-18, two presynaptic proteins, which are involved in sequential steps leading to vesicle exocytosis. Notably, these effect were strongly reduced in the presence of NaHCO $_3$, indicating that ONOO \cdot /ONOOH acts mainly intracellularly. Synaptophysin, one of the most abundant integral proteins of SV membrane, can be also nitrated (on Tyr250) and the formation of the synaptophysin/dynamin complex is impaired following ONOO \cdot /ONOOH exposure [114, 116]. Mallozzi et al. [114] have identified by LC-MS/MS analysis one major nitration site at Tyr354 in dynamin I isolated from synaptosomes treated with ONOO \cdot /ONOOH. LC-MS/MS analysis revealed also that in untreated synaptosomes dynamin I showed a basal level of nitration on Tyr125, Tyr541 and Tyr669; however, the low physiologic nitration level of these sites did not affect dynamin I functional properties. Instead, Tyr354 was nitrated only after ONOO \cdot /ONOOH treatment of synaptosomes implying that this site-specific post-translational modification likely accounts for dynamin I dysfunction. Vrljic *et al.* [117] detected nitration in 6 of 11 surface accessible Tyr residues of synaptotagmin 1 [three in the C2A domain (Tyr151, Tyr216 and Tyr229) and three in the C2B domain (Tyr311, Tyr364 and Tyr380)]. Synaptotagmin 1 is a Ca $^{2+}$ sensor for SNARE (soluble N-ethylmaleimide sensitive factor attachment protein receptor)-mediated, Ca $^{2+}$ -triggered synaptic vesicle fusion in neurons. Integration of the peak intensity for the individual synaptotagmin 1 peptides suggests a stoichiometry for the 3-NT modifications of 1–10% depending on the site, with the exception of Tyr151, which appears to be ~100% modified since an unmodified form was not identified [108].

Amyloid beta (A β) is a critical factor involved in the pathogenesis of AD. It was demonstrated that continuous intracerebroventricular infusion of A β 1–40 induced a time-dependent expression of the inducible nitric oxide synthase (iNOS) and an overproduction of NO in the rat hippocampus. The pathophysiological significance of the overproduction of NO for brain function was manifested by an impairment of nicotine-evoked acetylcholine (ACh) release and memory deficits [118]. Tran et al. [119] found that chronic A β 1–40 infusion caused a robust ONOO \cdot /ONOOH forma-

Table 3. Selected proteins nitrated in Alzheimer's disease.

Nitrated protein	Material	Methodology	Major observations	Ref.
Enolase	Male Wistar rat synaptosomes	WB	Nitration of enolase and synaptic proteins mediated by H ₂ O ₂ , NO ₂ and amyloid β heme peroxidase activity	[127]
Nitro-triosephosphate isomerase (nitration of tyrosines 164 and 208, close to the catalytic site)	Immunoprecipitates from hippocampus (9 individuals) and frontal cortex (13 individuals) of AD patients, compared with healthy subjects (4 and 9 individuals, respectively); Human embryonic kidney cells overexpressing mutant triosephosphate isomerase	WB, Transmission electron microscopy, Atomic force microscopy	Nitro-triosephosphate isomerase forms large beta-sheet aggregates <i>in vitro</i> and <i>in vivo</i> , Nitro-triosephosphate isomerase binds tau monomers and induces tau aggregation to form paired helical filaments	[128]
Brain proteins	Brain samples; normal control subjects: 4 females and 2 males, average age at death of 81 ± 6.4 y; amnesic mild cognitive impairment (MCI) patients, 4 females and 2 males, average age at death of 88 ± 3.8 y	Slot blot, Immunohistochemistry	Protein nitration is higher in the inferior parietal lobule (IPL) and hippocampus in MCI than in control subjects	[120]
α-Enolase, triosephosphate isomerase, neuropolypeptide h3, β-actin, L-lactate dehydrogenase, γ-enolase	IPL tissue specimens used for analyses taken at autopsy from five AD patients and five control subjects	WB, MS	Identification of six targets of protein nitration in AD suggests a role of protein modification by RNS in the progression of AD	[129]
α-Enolase, glyceraldehyde-3-phosphate dehydrogenase, ATP synthase alpha chain, carbonic anhydrase-II, voltage-dependent anion channel-protein (hippocampus)	Hippocampal samples from six AD patients and six age-matched controls	Immunoprecipitation, WB, MS	Nitration of proteins in AD hippocampus may be involved in the mechanisms of AD	[130]
Peroxiredoxin 2, triose phosphate isomerase, glutamate dehydrogenase, neuropolypeptide h3, phosphoglycerate mutase 1, H ⁺ -transporting ATPase, α-enolase, fructose-1,6-bisphosphate aldolase	IPL samples from four early AD (EAD) patients (79 ± 2 years) and four age-matched controls (average age at death of 86 ± 4 years).	WB, 2D PAGE, In-gel trypsin digestion, MS	The level of nitrated proteins in the IPL of early AD patients increased by 18% increase compared with age-matched controls	[131]

2D PAGE, two dimensional polyacrylamide gel electrophoresis; IPL, inferior parietal lobule; MS, mass spectrometry; WB, Western Blotting

tion and subsequent Tyr nitration of hippocampal proteins. Immunoprecipitation and Western blot analyses revealed that synaptophysin is a main target of Tyr nitration. These findings suggest that the Tyr nitration of synaptophysin is related to A β -induced impairment of ACh release.

Tyrosine nitration of brain proteins

Nitric oxide participates in the regulation of the daily activities of cells as well as in cytotoxic events. Tyr nitration is one specific form of protein modification that is associated with age-related neurodegenerative diseases [120]. Protein nitration enhances A β aggregation in a rodent model of AD [109]. By mediating Tyr nitration at the *ortho* position, ONOO⁻/ONOOH modification of proteins can block later phosphorylation events, thereby inducing protein dysfunction [121], some researchers have proposed that dynamic interplay between nitration and phosphorylation may be required for some normal biological functions, and Tyr nitration can contribute to differentiation of neuronal cell types and to neurite elongation.

The microtubule-associated tau protein is unfolded and finely soluble under physiological conditions, but in the brain tissue of AD changes in its conformation occur, affecting its solubility. Horiguchi et al. demonstrated the presence of nitrated tau protein in pretangles, neurofibrillary tangles, as well as tau inclusions in AD brain. Tau contains five Tyr residues (located at 18, 29, 197, 310, and 394), that can undergo nitration to initiate a range of 'tauopathies' [122]. In AD patients, the N-terminal tyrosine residues of the tau protein (Tyr18 and Tyr29) are more susceptible to nitrative modifications than other tyrosine residues (Tyr197 and Tyr394) [123]. Tau nitration at Tyr197 and Tyr18 has been reported to enhance disease progression in a range of neuro-degenerative disorders [124], whereas nitration at Tyr29 appears to be a specific characteristic of AD [123].

It was found that nitration of other proteins perturbs pH regulation, energy metabolism, and mitochondrial functions, and may be involved in the mechanisms of neuronal loss and progression of AD. In particular, such nitrated proteins were identified in the AD hippocampus as α -enolase, carbonic anhydrase II, glyceraldehyde-3-phosphate dehydrogenase, ATP synthase α -chain and voltage dependent anion channel protein 1 (VDAC-1), using a redox proteomics approach. Nitration of ATP synthase α -chain and VDAC-1 is associated with mitochondrial dysfunction and neuronal cell death in the AD hippocampus. Moreover, nitrated proteins are usually tagged for selective destruction in proteasomes, but in

AD this pathway may be defective due to oxidation of ubiquitin carboxy-terminal hydrolase L-1 in the inferior parietal lobule and hippocampus [122, 125, 126]. Main brain proteins nitrated in AD are listed in Table 3.

It should be noted that aging is an important risk factor for human α -synucleinopathies such as PD. There is a link between aging, α -synuclein (α Syn) abnormalities and enhanced vulnerability to neurodegenerative processes. It was also reported that phospho-Ser 129 and nitrated α Syn are formed within dopaminergic neurons of the monkey *substantia nigra* in the course of normal aging [132]. Schildknecht et al. [133] hypothesized that under physiological conditions α Syn may act as an intracellular scavenger of oxidants, catalytically regenerated, and performs an important protective role before the onset of disease or during aging. α Syn is a 140 amino-acid protein, originally identified in association with synaptic vesicles in the presynaptic nerve terminals and has been shown to interact with membranes both *in vitro* and *in vivo*. It is predominantly expressed in the brain (in the neocortex, hippocampus, *substantia nigra*, thalamus, and cerebellum, accounting for approximately 1% of brain weight) and is also present in other cells and tissues, including erythrocytes [134]. α Syn is involved in the modulation of synaptic activity through regulation of assembly of SNARE-complex of presynaptic vesicles, regulation of neurotransmitter release, regulation of cell differentiation and phospholipid metabolism [135]. Susceptibility to PD may be linked to modulation of α Syn protein expression. Furthermore, nitration of α Syn was associated with enhanced propensity of this protein to aggregate. Burai et al. [136] examined the site-specific incorporation of 3-NT at different regions of α Syn. They found that depending on the site of nitration, various nitrated α Syn species exhibit distinct structural and aggregation properties and exhibit reduced affinity to negatively charged vesicle membranes. Intermolecular interactions between the N- and C-terminal regions of α Syn play critical roles in mediating nitration-induced oligomerization of α Syn. In mutants, in which Tyr39 is not available for nitration, the extent of cross-linking is limited mostly to dimer formation, whereas mutants in which Tyr39 is available, along with one or multiple C-terminal tyrosines remain nitrated. Nitrated α Syn was observed to induce adaptive immune responses that exacerbate PD pathology in the mouse MPTP model of PD [137]. Increased nitrated α Syn is present in peripheral blood mononuclear cells of idiopathic PD patients compared to healthy individuals [138]. These studies provide evidence for a direct link between nitrative damage and the onset and progression of neurodegenerative synucleinopathies. More recently, Kleinknecht et al. [139] reported that Syn can be nitrated and form stable covalent dimers

originating from covalent crosslinking of two Tyr residues. Nitrated Tyr residues, but not dityrosine-crosslinked dimers, contribute to α Syn cytotoxicity and aggregation. Analysis of Tyr residues involved in nitration and crosslinking revealed that the C-terminus, rather than the N-terminus of α Syn, is modified by nitration and dityrosine formation. These data suggest that C-terminal Tyr133 plays a major role in α Syn aggregate clearance by supporting the protective Ser129 phosphorylation for autophagy and by promoting proteasomal clearance. C-terminal Tyr nitration increases pathogenicity and can only be partially detoxified by α Syn dityrosine dimers. It seems that complex interplay between Ser129 phosphorylation and C-terminal Tyr modifications of α Syn likely participates in PD pathology. Table 4 shows data on α -synuclein nitration in PD.

Nitration and aging

It has been extensively documented that increased nitration is often connected to the development of age-related diseases. High concentrations of peroxynitrous acid may affect modulation of mitochondrial respiration that can act as platform for development of prevalent neurodegenerative diseases. Proteomic ana-

lysis by ESI-MS/MS had shown that flotillin-1 and α -tubulin are nitrated in the rat in the course of aging. Age dependent accumulation of 3-NT on skeletal muscle glycogen phosphorylase b (Ph-b) is reported in an experimental rat model (106, 140). Results of selected studies on protein nitration in aging are shown in Table 5.

Methods used to measure the level of nitrated proteins

Among the many technologies available, the most effective and dependable method for the quantification of 3-NT are gas chromatography-mass spectrometry (GC-MS/MS) and liquid chromatography-mass spectrometry (LC-MS/MS). GC-MS/MS and LC-MS/MS based methods showed that the concentration of 3-NT in human plasma is on the threshold of the picomolar (pM) to nanomolar (nM) range and changes only very little upon disease or intervention. These important findings are suitable to serve as the gold standard and as a measure to test the reliability of alternative techniques, such as GC-MS, high performance liquid chromatography (HPLC) with electrochemical detection, or immunological assays.

Table 4. α -Synuclein nitration in Parkinson's disease.

Nitrated Protein	Materials	Methodology	Major observations	Reference
α Syn	Male Fischer 344 rats, 3-month-old vs 16-month-old	Western blotting, ELISA	Microglia activation and proinflammatory cytokine expression enhanced in the <i>substantia nigra</i> of elderly rats following intrapallidal lipopolysaccharide administration, Greater nitration of α Syn in the <i>substantia nigra</i> of 16-month-old rats vs 3-month-old rats, accompanied by a higher expression level of iNOS	[140]
α Syn	Twelve-month-old male nTg, SYN Tg, and SYN-null mice; primary neuronal and glial cultures.	WB, Immunostaining, Sequential biochemical fractionation, Immunoelectron microscopy	Neuroinflammation and Syn pathology are linked mechanistically to the onset and progression of PD	[141]
α -Syn (nitration of Tyr125 and Tyr136)	Squirrel monkeys of 2 age groups: <10 y (6–9 years, n=4) and >16 (17–19 y, n=3).	Immunohistochemistry	Age-related elevations of modified protein	[132]

Tg, transgene; WB, Western blotting

Table 5. Selected results of nitroproteomic studies of aging.

Material	Methods	Nitrated Protein	Findings	Reference
Male Wistar rats aged 6 m (adult) and 25 m (old)	WB	Quadriceps protein fractions	3-NT levels higher in all protein fractions of skeletal muscle in old male rats, especially in the mitochondrial fraction	[165]
Young (19–22 w) and old (24 m) C57BL//6 male mice	SDS PAGE, WB	Hepatic proteins	Significantly higher level of Tyr nitration of proteins in old mice vs young mice	[166]
PC12 cell culture	WB, MS	Actin, tubulin, Hsp70, Hsp90	ONOOH-treated Hsp70, actin, and tubulin nontoxic for motor neurons and PC12 cells. ONOOH-treated Hsp90 induced death in ~40% of PC12 cells and 60% of motor neurons	[167]
Young adult (4-5 m), middle-aged (10 and 16 m) and old (26-28 m) Fisher 344 male rats	WB, HPLC-MS	SERCA2a nitrated at Tyr294 and Tyr295	Age-dependent nitration and loss of function of the rat skeletal-muscle SR Ca ²⁺ -ATPase isoforms SERCA1 and SERCA2a	[168]
18 male F344 rats were 7–11 m old (young adult), 22–25 m old (old), and 27–30 m old (very old)	WB	SERCA2a, aconitase, β-enolase, carbonic anhydrase III, triosephosphate isomerase	Significant age-associated increase in nitrotyrosine-modified proteins	[169]
Male F344 BN/F1 rats aged 5, 22, and 34 m	WB, MS/MS	LDL receptor related protein 2, CNP and others	Age-dependent accumulation of nitrated proteins	[170]
Young (4 m) and old (24 m) Fisher344 rats and young (6 m) and old (34 m) Fisher 344 /BN F1 rats	WB, MALDI-TOF MS	α-Fructose aldolase, triosephosphate isomerase, GAPDH and others	Nitrated proteins accumulate at a faster rate in old compared to young tissue, Nitrated proteins are subject to proteasomal degradation, Proteasomal activity declines with increasing age	[171]
17 Fisher 344/BN F1 rats (10-34 m old and 7-5 m old)	WB, MS/MS	Tropomyosin 1 - α isoform, neurofibromin, cadherin EGF-LAG, seven pass G type receptor 2	Nitrated proteins present in cardiac tissue, their abundance increases with age, 1.5 to 2 fold increase in protein nitration in 34-m vs 5-m old animals	[172]
Young (4-6 m old) and aged (24-26 m old) male C57BL/6 mice	WB, MALDI TOF-MS	Profilin 1, polymerase I, Transcript release factor, peroxiredoxin 6, and others	Significant modification of vascular endothelial cytoskeleton, which potentially contributes to barrier dysfunction, increased vascular permeability and pulmonary oedema	[173]

CNP, 2,3-cyclic nucleotide 3-phosphodiesterase; MS, mass spectrometry; WB, Western blotting

The various antibody assays also need to be validated by these GC–MS/MS or LC–MS/MS methods [141]. Quantitative MS-based analysis is essential for the elucidation of the stoichiometry of the specific tau Lys-directed posttranslation protein modifications that correlate with AD neuropathology. Multiple Reaction Monitoring (MRM) is a targeted mass spectrometry (MS)-based technology that is becoming increasingly utilized for protein quantification. MRM-based approaches have been used to determine the relative abundance of tau polyubiquitylation in human AD brain and global tau in human CSF. In contrast to MS-based discovery proteomics experiments, MRM entails the targeted, simultaneous measurements of peptides that serve as surrogates for the protein targets of interest. MRM-based assays are considered to be the “gold standard” for MS-based targeted protein quantification since they are highly specific, precise, and accurate, and they can be multiplexed (hundreds of peptides can be quantified in a single assay), standardized and readily reproduced. A targeted proteomics method that is similar to MRM is parallel reaction monitoring (PRM) wherein an accurate mass and high-resolution mass spectrometer is used to permit the parallel detection of all target product ions [142].

CHLORINATIVE PROTEIN MODIFICATIONS IN AGING AND AGE-RELATED NEURODEGENERATIVE DISEASES

Chlorinative stress undoubtedly contributes to the pathogenesis of neurodegenerative diseases [143]. In brain, chloride ions are present at the concentration of $10^{-2} - 10^{-1}$ M [144]. HOCl can be generated with the activation of microglia and myeloperoxidase secretion [145-148]. Moreover, infiltration of monocyte/macrophage and neuronal expression of myeloperoxidase also contribute to the formation of HOCl [149, 150]. Suppo-

sedly, the brain has poor defence system against HOCl [79,151,152]. Thus, the toxicity of HOCl towards central nervous system tissue was shown [153-155]. Furthermore, MPO was reported to be expressed with increased levels in the cerebral tissue of patients affected by AD [156] and 3-chlorotyrosine as a biomarker of HOCl production was detected in proteins from AD hippocampus. The level of 3-chlorotyrosine in the samples from diseased brain was three-fold higher compared to control samples [150]. Halogenation has a clear effect on the self-assembly of the amyloid β peptide aggregates [157]. However, it can be concluded that a role of protein chlorination in neurodegenerative diseases is not analysed completely yet.

ADVANCED PROTEIN OXIDATION PRODUCTS (AOPP) IN AGING AND AGE-RELATED DISEASE

A special class of protein modification products, consisting of oxidized, dityrosine-containing, crosslinked proteins formed mainly by reactions of RXS with plasma proteins, predominantly albumin, are so-called advanced oxidation protein products (AOPP). *In vivo*, the generation of chlorinated oxidants is a feature of phagocytic cells containing MPO [158, 159]. Witko-Sarsat et al. [160] first reported elevated plasma level of AOPPs in uremic patients. High levels of AOPPs were detected in patients on maintenance hemodialysis, followed by those on peritoneal dialysis. Patients with advanced chronic renal failure not yet on dialysis had almost three times higher AOPP levels than healthy subjects.

Size exclusion chromatography of uremic plasma has isolated high-molecular-weight (600 kDa) and low-molecular-weight (80 kDa) AOPPs. The high molecular-weight AOPPs were mostly formed of albumin aggregates, likely resulting from disulfide bridges and/or

Table 6. Chosen results of studies on the effect of aging and neurodegenerative diseases on the AOPP level in blood serum or plasma.

Subjects studied	AOPP level	Reference
Alzheimer disease	Increased (106.5±27.3 vs 87.5±37.8 μ M)	[161]
Chronic schizophrenia	Increased (211.2±159.4 vs 191.7± 146.3 μ M)	[162]
Parkinson disease	Increased (65.6 vs 45.6 μ M)	[163]
Postmenopausal vs premenopausal women	Increased (118.6±59.1 vs 61.6 ± 16.4 μ M)	[164]
Systemic sclerosis	Increased (109.1 vs 75.5 μ M)	[165]
Rats, 9-m old (adult) vs 3-m old (young)	Increased (8.3±2.7 vs 6.8±2.3 μ M)	[166]
Rats, 22-m old (old) vs 3-m old (young)	Increased (16.1± 4.8 vs 6.8±2.3 μ M)	[167]
Rats, 22-m old vs 2-m old	Increased (198.5±44.9 vs 129.3±27.2 μ M)	[168]

dityrosine crosslinking. The low molecular weight of AOPPs contained albumin in the monomeric form [160].

Exemplary values of AOPP obtained in studies of aging and neurodegenerative diseases are reported in Table 6.

AOPPs were first recognized as markers of oxidative stress. However, it was reported that AOPPs can also promote ROS production, which leads to a vicious circle. AOPPs activate NADPH oxidase via the protein kinase C-dependent pathway inducing an excessive generation of intracellular superoxide in various renal cells (podocytes, endothelial cells, mesangial cells, and tubular epithelial cells) [169].

AOPPs are assayed spectrophotometrically at 340 nm after treatment samples with KI [160, 170]. A kinetic AOPP assay has been proposed [171], but a limited correlation was found to exist between results obtained by the classical and kinetic assay [172].

BRAIN PROTEIN MODIFICATIONS BY 4-HYDROXY-2,3-TRANS-NONENAL IN AGING AND NEURODEGENERATIVE DISEASES

Post-mitotic neurons are notably vulnerable to lipid peroxidation since the brain has high levels of polyunsaturated fatty acids, high levels of redox transition metal ions, high oxygen consumption, relatively low

levels of low-molecular weight anti-oxidants and antioxidant enzymes. Peroxidation of polyunsaturated fatty acids, especially linoleic acid, linolenic acid and arachidonic acid by non-enzymatic processes leads to the formation of aldehydes, among them 4-HNE is present at very low concentration in plasma, in the range of 0.28–0.68 μM under physiologic conditions, but its concentration in cells, where it is produced, may be higher ($\leq 5 \mu\text{M}$) [173]. 4-HNE concentration can be increased as much as by 100 times under OS conditions [174]. Esterbauer's group demonstrated that 4-HNE formation from arachidonic acid is greater in the presence of NADPH-dependent microsomal enzymes [175]. 4-HNE possesses three reactive functions: a C2=C3 double bond, a C1=O carbonyl group and a hydroxyl group on C4. These functions make this electrophilic molecule highly reactive toward nucleophilic thiol and amino groups. 4-HNE can enter the reaction of Michael addition to thiol or amino groups, which involves the C3 of the C2=C3 double bond or can form Schiff bases between the C1 carbonyl group and primary amines. The kinetics of the Schiff base formation is slow and reversible, making Michael-adducts predominant adducts of 4-HNE to proteins. 4-HNE reacts mainly His, Cys and Lys residues in proteins [176, 177] (Fig. 1F, Fig. 2). The formation of the 4-HNE-protein adducts is a bio-active marker of pathophysiological processes [178-180]. 4-HNE forms Michael adducts with enzyme peptidylprolyl cis/trans-isomerase A1 (Pin1), which catalyzes conversions of




Figure 2. Reactions of 4-hydroxy-2,3-*trans*-nonenal (4-HNE) with proteins.

phospho-serine and phospho-threonine-proline from *cis* to *trans* conformation. These adducts were detected by matrix-assisted laser desorption/ionization/time-of-flight/time-of-flight (MALDI-TOF/TOF) mass spectrometry at the active site residues His157 and Cys113, with Cys113 being the primary site of 4-HNE modification [181-185]. Protein modifications by 4-HNE impairs glutamate and glucose transport, disrupts Ca^{2+} homeostasis, damages cholinergic neurons thus impairing visuospatial memory and induces apoptosis in PC12 cells (cell line derived from a pheochromocytoma of the rat adrenal medulla) and cultured rat hippocampal neurons [186-188]. Nam et al. (2014) compared N-methyl-D-aspartate receptor type 1 (NMDAR1) and 4-HNE in the hippocampus of D-galactose (D-gal)-induced and naturally aging models of mice [189]. These authors observed an age-dependent reduction of NMDAR1 and an increase in 4-HNE in the dentate gyrus, CA1 and CA3 regions of the hippocampus *via* immunohistochemistry and Western blot analyses. In the D-gal-induced chemical aging model they noted similar changes in NMDAR1 and 4-HNE although the degree of reduction/increase in NMDAR1/HNE was not as severe as that in the naturally aging mice.

4-HNE-protein adducts were found to be elevated in brain tissues and body fluids of AD, PD, Huntington disease as well as ALS subjects [190, 191]. 4-HNE-His adducts were reactive with A β core of senile plaques and neurofibrillary tangles [179]. Hardas et al. (2013) detected oxidative modification of lipoic acid, a key cofactor for a number of proteins including pyruvate dehydrogenase and α -ketoglutarate dehydrogenase, by 4-HNE in AD brain [192]. In another study, 4-HNE-Lys adducts were increased in neurons containing neurofibrillary tangles, but also in pyramidal neurons located in the hippocampal tissue sections in AD [193]. The formation 4-HNE adducts with the neuronal glucose transporter GLUT3 and the mitochondrial ATP synthase α subunit in AD brain leads to reduced glucose utilization and energy production in AD [194, 195]. Studies conducted by Sultana et al. suggest that 4-HNE-modification of α -enolase, heme oxygenase 1, Collapsin Response Mediator Protein-2 and ATP synthase subunit α are critical in the progression of AD [196]. These authors hypothesized that 4-HNE modification can be not a random event, but occurs on specific proteins, which, in turn, display altered functions. The formation of 4-HNE adducts with α -enolase could inhibit the conversion of plasminogen to plasmin and the degradation of A β . In AD brains, the increase of OS leads also to increases of Nrf2 activity as well as, consequently, increases of heme oxygenase 1 level. Heme oxygenase 1 catalyzes the degradation of heme and represents the rate-limiting enzyme in bilirubin production [197]. Collapsin Response Mediator Protein-

2 (dihydropyrimidinase-related protein-2) plays an important role in cytoskeletal organization, axonogenesis, axon outgrowth, membrane trafficking and neuronal polarity [198]. The oxidative modification of Collapsin Response Mediator Protein-2, such as formation adducts with 4-HNE, can play an important role in shortening of axons as well as loss of synapses in AD. ATP synthase subunit α , a part of complex V responsible for mitochondrial-resident ATP synthesis. ATP synthase α might be modified by 4-HNE in AD brain, which causes the reduced activity of ATP synthase and reduced ATP levels in AD brain compared to age-matched controls [196]. According to recent study, klotho gene therapy in senescence-accelerated mouse prone-8 (SAMP8) reduced memory deficits, neuronal loss, synaptic damage and 4-HNE levels, and increased mitochondrial SOD-2 and catalase expression. Additionally, the up-regulation of klotho expression decreased Akt and Forkhead box class O1 (FoxO1) phosphorylation. The role of 4-HNE adducts in ALS progression has been recently reviewed by Zarkovic group [180]. ALS is a progressive neurodegenerative disorder characterized by weakness and spasticity, caused by the loss of lower and of upper motor neurons and by secondary neurogenic amyotrophy of striated muscles. An *in vitro* study demonstrated that 4-HNE impairs the glutamate and glucose transport and the choline acetyltransferase activity in cultured motor neurons [199], while human autopsy materials have shown increased levels of 4-HNE, which modifies astrocytic glutamate transporter EAAT2 (excitatory amino acid transporter 2) impairing glutamate transport in ALS. Moreover, 4-HNE is able to target SOD1 in ALS [200]. Kabuta et al. (2015) reported that TDP-43, a major component of ubiquitin-positive inclusions in ALS, is bound by 4-HNE, therefore inducing both proteins into toxic aggregates [201].

It should be mentioned that 4-HNE has also crucial role in α Syn-induced cytotoxicity and neuro-inflammation [202]. These aldehydes can also promote the formation of α Syn oligomers with defined structural properties. Although, 4-HNE modifies α Syn immediately, primarily the His50 residue, oligomer formation only occurs with prolonged incubation times (> 24 h) and involving fewer cross-linking events. 4-HNE can bind to α Syn at an acidic pH, but these modifications cannot promote oligomerization even with increased incubation times [203]. The current objective of research in the field of contribution of 4-HNE-protein adducts is characterization the interactions of 4-HNE with redox sensitive cell signalling proteins. 4-HNE is involved in aging-related signaling pathways, such as NF- κ B, AKT, Nrf2 and mTOR. Other signaling pathways involved in aging, for example related to growth factor signaling EGFR, PDGFR and others are also modified by 4-HNE.

Understanding how modulation of activities of these signaling pathways contributes to physiological aging and neurodegenerative diseases may pave the way for new therapeutic strategies.

Assay of 4-HNE-protein adducts

The gold standard in studies of protein modifications by lipid peroxidation products, including 4-HNE, in proteomic studies is mass spectrometry, e. g. matrix-assisted laser desorption ionization/time-of-flight/time-of-flight (MALDI-TOF/TOF), ESI-MS or LC-ESI-CID-MS/MS [204-206]. Antibodies against the His adduct of 4-HNE has allowed for facile detection and quantification of 4-HNE-modified proteins by immunochemical techniques (immunoblotting, immunocytochemistry, immunohistochemistry and immuno-electron microscopy).

Two variants of the 4-HNE-ELISA assay have been developed, both of which are based on the 4-HNE-His monoclonal antibodies. The differences between these two assays concern the analytical protocols and the albumin-HNE standards used, allowing very sensitive determination of low amounts of the 4-HNE-protein adducts (the assay denoted *HNE-His ELISA Fine*) even below 0.025 nmol 4-HNE-His/mg of protein and the one able to detect higher amounts, above 1.5 nmol 4-HNE-His/ mg of protein (the assay denoted *HNE-His ELISA Stress*) [207].

ROLE OF OXITATIVE STRESS IN THE BLOOD BRAIN BARRIER AGING

The blood brain barrier (BBB) separates the brain and blood with a large surface area (between 12 and 18 m² in the average human adult) [208, 209]. The opposing membranes of endothelial cells are connected by tight junctions, which are formed through an intricate network of interacting proteins such as claudins, occludin, junctional adhesion molecules and cytoplasmic proteins [210]. Nitta et al. (2003) demonstrated that claudin-5 is a critical determinant of BBB permeability [211]. In the process of healthy aging an increased “leakage” of BBB may occur, not only due to alteration of thickness of basal lamina, endothelial cells, morphology of pericytes and astrocytes, but also as a result of the changes in expression of transporter proteins at the endothelial cell layer of BBB [212]. Bors et al. (2018) reported that the number of tight junctions decreases, the thickness of basal lamina increases as well as the size of astrocyte endfeet extends with advanced age. These authors also demonstrated that the function of P-glycoprotein 1 (P-gp, ABCB1 Abcb1a/Mdr1a), the most important efflux transporter located on the luminal surface of brain capillary

endothelial cells is reduced in old Wistar rats [213]. Reduced BBB expression of P-gp was associated with increased brain parenchymal A β 40 and A β 42 levels in aged rats [214], in agreement with the idea that P-gp is an important efflux transporter to remove A β from the CNS [215]. Pan et al. (2018) showed that low density lipoprotein receptor-related protein 1 (LRP-1) expression declines with age, which may contribute to A β accumulation [209]. Van Assema et al. (2012) studied *in vivo* effects of gender and aging on human BBB P-gp function in a large sample size using PET and (R)-[¹¹C]verapamil. These authors reported that decreased BBB P-gp is found with aging; nevertheless, effects of age on BBB P-gp function differ between men and women [216].

The function of BBB can be impaired by ROS/RNS, and these effects are partly mediated by products of lipid peroxidation [217]. The major secondary lipid peroxidation product, 4-HNE can impair the BBB function *via* the decrease of GSH [218]. Wang et al. (2012) reported that overexpression of actin-depolymerizing factor (ADF) blocks the oxidized low-density lipoprotein (ox-LDL)-induced disruption of endothelial barrier. Furthermore, siRNA-mediated downregulation of ADF expression aggravated ox-LDL-induced disruption of endothelial barrier and ROS formation. ADF seems to be a key signaling molecule in the regulation of BBB integrity and suggest that ADF might be used as a target to modulate diseases accompanied by ox-LDL-induced BBB compromise [219]. It should be also mentioned that several studies suggest a link between synucleinopathies and the cholesterol metabolite 27-hydroxycholesterol (27-OHC). 27-OHC is the major cholesterol metabolite in the blood that crosses BBB, and its levels can increase following hypercholesterolemia, aging and OS, which are all factors for increased synucleinopathy risk. 27-OHC can increase α Syn levels and causes the inhibition of the proteasomal function and reduction in heat shock protein 70 levels as potential cellular mechanisms involved in regulation of α Syn [220].

REMOVAL OF MODIFIED PROTEINS

The level of posttranslationally modified proteins is a resultant of the rate of protein modification and rate of removal of modified proteins. Aging, as well as several age-related diseases are associated with a decreased ability to maintain proteostasis [221]. All cells have a number of quality control mechanisms in order to maintain the stability and functionality of their proteome. The proteostasis network includes both protein stabilization mechanisms (major heat shock proteins) and protein degradation systems (proteasome and lysosome) [222-224]. In addition, there are several

modulators of proteotoxicity (like MOAG-4), that operate through distinct pathways [42]. All these systems work in concert to restore the structure of denatured proteins or to promote their degradation, thus preventing the accumulation of damaged components and ensuring the continuous renewal of the intracellular polypeptides. Many studies have shown that aging is accompanied by failure of proteostasis [225], while chronic exposure to denatured or aggregated proteins contributes to the development of age-related neuro-degenerative diseases such as AD and PD [221, 226].

The proteasome

The proteasome is a fundamental multicatalytic enzyme complex, which facilitates the degradation of normal as well as abnormal, damaged, denatured and redundant cellular proteins. Proteasomes are located in different cellular compartments (cytoplasm, nucleus and endoplasmic reticulum) and represent approximately up to 1% of the total cellular protein content. The central role of the proteasomes is demonstrated by their participation in numerous and diverse cellular functions, including the regulation of transcription factor abundance, cell cycle and cellular differentiation. The main proteasomal complex is the 30S/26S proteasome and is composed by the 20S catalytic "core" and the 19S regulatory "cap" (summarized in [227]).

The 20S proteasome is a barrel-like structure composed of 28 protein subunits that form a complex of 700 kDa. The two outer rings comprise seven different α subunits, while the interior rings consist of seven β subunits, creating an $\alpha 1-7/\beta 1-7/\beta 1-7/\alpha 1-7$ layout. The external α rings control the entry of proteasome's substrates into the β rings, the site of the proteolytic activity. The α -subunits are additionally responsible for the binding of different factors that regulate the activity and specificity of the catalytic core. Three of the seven β subunits, namely $\beta 1$, $\beta 2$ and $\beta 5$, are proteolytically active, having different substrate specificity. Specifically, $\beta 1$ has a caspase-like activity (CL or PGPH), $\beta 2$ a trypsin-like (TL) and $\beta 5$ a chymotrypsin-like activity (CT-L). The protein hydrolysis occurs after acidic peptide bonds, basic amino acids and hydrophobic amino acids, respectively [228].

The 19S regulatory complex is composed of 19 different subunits that form two heteromeric rings, known as "base" and "lid" [182]. It is responsible for binding, deubiquitination and translocation of the protein substrate in the 20S core. The base is composed of nine subunits, 6 of which (Rpt1-6) possess ATPase activity [230]. Rpn1, Rpn2 and Rpn13 are 3 non-

ATPases that are necessary for the proper function of the 19S complex. In addition, since they act as polyubiquitin receptors, these subunits are responsible for the recognition of the ubiquitinated protein substrate [231]. The "lid" bridges the gap between the 20S and the 19S proteasomal particles. This structure is evolutionary conserved and consists of nine RPN subunits (Rpn3, 5 -9, 11, 12 and 15). The "lid" is very flexible structure, necessary for the positioning and the deubiquitination of the substrate by the deubiquitinating subunit Rpn11 [232]. Thus, the 19S regulatory complex acts as a very versatile device, which facilitates the access of the protein substrate to the core of the 20S proteasome in an ATP-dependent manner.

The 26S/30S proteasome is formed by the 20S catalytic core and the 19S regulatory particle. One or two regulatory complexes may bind on the catalytic core, forming the 26S or the 30S complexes, respectively. The substrates of the 26S proteasome are identified by labeling with multiple ubiquitin molecules. The ubiquitin is attached via a three-step procedure, which requires the action of E1 (ubiquitin activation), E2 (ubiquitin conjugation) and E3 (ubiquitin ligase) ligases. Polymeric ubiquitin chains are produced by the repeated action of the E1, E2 and E3 enzymes. The multi-ubiquitin chains signal the identification of the protein substrate for degradation. Upon recognition of the substrate, the poly-ubiquitin chains are removed by deubiquitinating enzymes (DUBs) [226]. The overall mechanisms of ubiquitination and proteasomal degradation are known as the *ubiquitin-proteasome system* (UPS system) (Fig. 3).

Besides the constitutive proteasomes, there are specific specialized proteasomes, formed when the $\beta 1$, $\beta 2$ and $\beta 5$ catalytic subunits become *de novo* substituted by $\beta 1i$, $\beta 2i$ and $\beta 5i$ subunits, respectively. These subunits are induced in response to the immunomodulatory cytokine interferon-gamma (IFN-gamma). The immunoproteasomes, as they are termed, besides their main role in antigen presentation, are involved in adaptation to OS and in selective degradation of oxidized proteins during aging, possibly in response to chronic inflammation (as summarized in [226]).

Proteasome and aging

During aging proteostasis collapses [223], resulting in the accumulation of denatured, aggregated or oxidized proteins, which in turn causes cellular damage and impairment of tissues [233]. The proteasomes, being the main proteolytic cellular system responsible for the elimination of nonfunctional or excessive proteins, hold a pivotal role in aging [234].




Figure 3. Overview of the ubiquitin (Ub)/proteasome system and its substrates in relation to aging. Ub conjugation is mediated by a series of enzymes. The Ub-activating enzyme E1 transfers Ub to the active site of the E2 Ub-conjugating enzyme and the E3 Ub-ligase ligate Ub to the target protein. The ubiquitinated protein is targeted to the 26S proteasome for degradation. The 26S proteasome consists of the 20S catalytic core and of one or two 19S regulatory particles. The 20S proteasome consists of 28 subunits that are divided to two outer α and two central β rings. The immunoproteasome is induced in response to the immunomodulatory cytokine interferon-gamma (IFN-gamma) or in response to the increased OS that is observed during aging. The age-related elevation of OS also causes oxidative damage to proteins, such as carbonylation. In addition, the excessive \bullet NO production during aging can lead to aberrant S-nitrosylation/tyrosine nitration. Nitrate proteins are prone to aggregation and may contribute to the onset and progression of various neurodegenerative diseases, including AD or PD. The accumulation of aggregated or carbonylated proteins inhibits proteasomal activity contributing to the observed proteasomal dysfunction during aging and to the advancement of age-related pathologies.

Young cells and organisms are characterized by an effective preservation of proteostasis. However, this ability is reduced during normal aging. This is evidenced by the increased accumulation of oxidatively modified proteins in senescent cells and tissues, which is indicative of the impairment of protein quality control and of protein degradation systems. Senescent cells have higher levels of proteins bearing modifications, such as oxidative carbonylation, oxidized Met and glycation. Studies *in vivo* and *in vitro* have shown that both the expression and function of the proteasome

are negatively affected by aging. Proteasome dysfunction during aging results not only due to the reduced expression of proteasome subunits and the impaired assembly of proteasomal complexes, but also because of the aggregated proteins that inhibit its function. Specifically, the reduction of proteasome activity during aging has been detected in numerous aged human tissues (muscles, lenses, skin, lymphocytes) or other mammalian tissues/organs such as the heart, muscles, spine, brain, liver, adipose tissue and retina (reviewed in [235]).

The activities of the proteasomes decline in senescent human fibroblasts, as a result of a reduction in expression of β subunits [236]. Moreover, it has been shown that the partial inhibition of the proteasomes in young cells causes a p53-mediated premature senescence [237]. On the other hand, the accumulation of damaged proteinaceous materials such as lipofuscin [238] or of protein aggregates [239] during aging, impairs proteasome function. Furthermore, studies in *D. melanogaster* have shown that the age-related disturbances of the 26S proteasome assembly lead to decreased proteasomal activity [240, 241]. Notably the naked mole, which is an extremely long-lived rodent, has high levels of proteasome activity, which may contribute to proteostasis maintenance and consequently to the extremely increased lifespan of these animals [242]. Similarly, fibroblasts derived from healthy centenarians have functional proteasomes, with characteristics similar to those of proteasomes from younger donors [243]. Accordingly, human embryonic stem cells (hESCs), that have an unlimited proliferative capacity, exhibit high proteasome activities, as compared to their differentiated counterparts [244]. Recently, the age-related decline of proteasome content and activities, along with the altered proteasome assembly, has been linked with the senescence-related loss of hMSC stemness [245]. Collectively, these studies demonstrate that aging is tightly connected with failures in biosynthesis, assembly and function of the proteasome.

Proteasome activation

Proteostasis failure is an important determinant of the aging process and is caused by a progressive decline of the respective defense systems. As such, interventions that promote proteostasis may delay aging and reduce the incidence of age-related diseases [246]. For instance, the activation of epidermal growth factor (EGF) signaling extends longevity in nematodes, by increasing the expression of various components of the ubiquitin-proteasome system [247]. Likewise, the enhancement of proteasome activity by deubiquitination inhibitors or by proteasome activators increases the replicative lifespan of yeast *Saccharomyces cerevisiae* [248]. In addition, the overexpression of the $\beta 5$ catalytic subunit [228] or of the 19S subunit Rpn6 [249] confers an increased lifespan in *C. elegans*.

Similar approaches for activating proteasomes have also proved successful in mammals. The genetic activation of the proteasome has been achieved by the stable overexpression of the catalytic $\beta 5$ subunit in the fibroblast cell lines WI-38/T and IMR90 [236]. These transfectants have increased ability to degrade oxidized proteins effectively, improved resistance to OS, while

the primary IMR90 cells display a 15-20% prolongation of their lifespan. Similarly, the restoration of normal levels of catalytic proteasome subunits ameliorates the aging phenotype in fibroblasts from elderly donors [250]. Overexpression of $\beta 5$ also promotes proteolysis and resistance to oxidative stress in human epithelial cells [251] and in promyelocytic leukemia HL60 cells [236]. Similar data have been reported in other cell types using different proteasome subunits. For instance, the overexpression of $\beta 6$ in human bronchial epithelial Beas2B cells increases the activity of the proteasome and protects against the endoplasmic reticulum (ER) stress induced by cigarette smoke [252]. Moreover, an elevation in expression levels of hUMP1/POMP, a chaperone facilitating proteasome assembly, results in increased proteasome activity and protects the cells from OS [205]. Similarly, an increase of PA28 levels in mouse cardiomyocytes stimulates the degradation of denatured proteins, protecting from heart proteinopathy [254]. Additionally, the overexpression of the regulatory 19S subunit Rpn6/PSMD11 enhances the assembly of 26S proteasome in human embryonic stem cells (hESCs) [243]. Remarkably, it has been recently revealed that overexpression of the $\beta 5$ proteasome subunit in human Wharton-Jelly derived mesenchymal stem cells (WJ-MSCs) resulted not only in increased proteasome activity and assembly, but also induced the expression of additional 26S proteasome subunits. The enhanced proteasome activity was maintained even after extensive culture, protecting the stem cells from the age-related increase of oxidative damage, as indicated by the reduced levels of ROS and of oxidatively modified proteins. Importantly, proteasome activation doubled the replicative lifespan, improved the expression of the core pluripotency factors and enhanced the differentiation capability towards adipocytes, osteocytes and chondrocytes of both young and senescent WJ-MSCs [245].

As genetic manipulation is not always feasible for clinical applications, there has been an effort towards the identification of natural or synthetic proteasome activators with antioxidant and anti-aging properties. Substances that directly induce the activity of the proteasome include pollen [255, 256], oleuropein [256], curcumin [258] and the synthetic peptide PAPI (Proteasome Activating Peptide-1) [259]. A different approach concerns the use of compounds that activate the transcription of proteasomal subunits. It is known that the transcription factor Nrf2 (Nuclear factor (erythroid-derived 2)-like 2) induces the expression of antioxidant enzymes including proteasomal subunits [260]. Treatment with 18 α -glycyrrhetic acid (18 α -GA) activates Nrf2, which in turn induces proteasome function and results in an enhancement of lifespan of both human fibroblasts [261] and *C. elegans* nematodes

[262]. Likewise, treatment with quercetin increases the CT-L proteasomal activity of human fibroblasts and increases their resistance to OS [263]. Finally, activation of Nrf2 by sulforaphane increases pluri-potency and self-renewal capacity of hESCs [264]. The analysis of the role of proteostasis maintenance mechanisms in aging, is essential for the rational design of interventions to improve the quality of human life in old age ('healthspan'), including the treatment of age-related diseases.

PERSPECTIVES

Abundant evidence demonstrates accumulation of products of protein modifications by ROS, RNS and RXS during aging of humans and model organisms and enhanced accumulation of such products in age-related diseases. New methods of analysis, based mainly on the MS technique, became available allowing for more precise identification of protein modifications and perhaps introduction of specific disease markers. Elucidation of the role of such modifications in aging-related changes and in the progress of diseases is more difficult. Are they only markers of aging and diseases or play a primary role in their development? There are reasons to not exclude the second possibility as these modifications adversely affect protein functions and interactions. Prospective and intervention studies may be helpful in this respect and may point to the possible role of specific protein modifications as possible early disease markers.

Abbreviations

2D PAGE: two-dimensional polyacrylamide gel electrophoresis; 3-NT: 3-nitrotyrosine; A β , Amyloid beta; ACR: acrolein; AD: Alzheimer's disease; ALS: amyotrophic lateral sclerosis; AOPP: Advanced Oxidation Protein Products; DNP: dinitrophenyl; DNPH: dinitrophenylhydrazine; ECM: extracellular matrix; ESI: electro spray ionization; GC: gas chromatography; 4-HNE: 4-Hydroxy-2,3-*trans*-nonenal; IP: immunoprecipitation; IPL: inferior parietal lobule; MDA: malondialdehyde; MCI: mild cognitive impairment; MPO: myeloperoxidase; MPTP: 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; MRM: Multiple Reaction Monitoring; MS: mass spectrometry; NOS: nitric oxide synthase; NS: nitr(os)ative stress; OS, oxidative stress; PD, Parkinson's disease; RCS, reactive carbonyl species; RNS, reactive nitrogen species; ROS: reactive oxygen species; RXS: reactive halogen species; SERCA: sarcoplasmic reticulum Ca²⁺-ATPase; SOD: superoxide dismutase; α Syn: α -synuclein; Tg: transgene; WB: Western blot; XS: halogenative stress.

AUTHOR CONTRIBUTIONS

Efstathios S. Gonos and Mariana Kapetanou are responsible for description of the removal of modified proteins, Jolanta Sereikaite for description of chlorinative protein modifications, Izabela Sadowska-Bartosz for description of the remaining part of the manuscript and general edition; Katarzyna Naparło and Michalina Grzesik provided part of data concerning protein carbonylation and nitration. Grzegorz Bartosz contributed to correction of the manuscript.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

FUNDING

This study was performed within the project „Nanomolecular antioxidants: biological basis of targeted therapy of neurodegenerative diseases” (number 2016/22/E/NZ7/00641) financed by National Science Centre (NCN), Poland in a programme „SONATA-BIS 6”. The paper is also a result of realization of research projects OPUS 9 (number 2015/17/B/NZ3/03731) financed by the NCN, Poland.

REFERENCES

1. Global Health and Aging - World Health Organization. National Institute on Aging National Institutes of Health NIH Publication. 2011; 11-7737.
2. World Population Ageing. 1950–2050. Department of Economic and Social Affairs. Population Division. United Nations. 2001.
3. Mikkelsen RB, Wardman P. Biological chemistry of reactive oxygen and nitrogen and radiation-induced signal transduction mechanisms. *Oncogene*. 2003; 22:5734–54. <https://doi.org/10.1038/sj.onc.1206663>
4. Sadowska-Bartosz I, Ott C, Grune T, Bartosz G. Posttranslational protein modifications by reactive nitrogen and chlorine species and strategies for their prevention and elimination. *Free Radic Res*. 2014; 48:1267–84. <https://doi.org/10.3109/10715762.2014.953494>
5. Bartosz G. Reactive oxygen species: destroyers or messengers? *Biochem Pharmacol*. 2009; 77:1303–15. <https://doi.org/10.1016/j.bcp.2008.11.009>
6. Barbieri E, Sestili P. Reactive oxygen species in skeletal muscle signaling. *J Signal Transduct*. 2012; 2012:982794. <https://doi.org/10.1155/2012/982794>
7. Sadowska-Bartosz I, Bartosz G. Chapter 1: Oxidative

- nitritative and chlorinative stress: biomarkers, in: *Studies on Psychiatric Disorders. Series: Oxidative Stress in Applied Basic Research and Clinical Practice.* Humana Press. 2015; XVIII:1-39.
8. Jung T, Höhn A, Grune T. The proteasome and the degradation of oxidized proteins: part II - protein oxidation and proteasomal degradation. *Redox Biol.* 2014; 2:99–104. <https://doi.org/10.1016/j.redox.2013.12.008>
 9. Davies MJ. Protein oxidation and peroxidation. *Biochem J.* 2016; 473:805–25. <https://doi.org/10.1042/BJ20151227>
 10. Gebicki JM. Protein hydroperoxides as new reactive oxygen species. *Redox Rep.* 1997; 3:99–110. <https://doi.org/10.1080/13510002.1997.11747096>
 11. Dalle-Donne I, Rossi R, Giustarini D, Milzani A, Colombo R. Protein carbonyl groups as biomarkers of oxidative stress. *Clin Chim Acta.* 2003; 329:23–38. [https://doi.org/10.1016/S0009-8981\(03\)00003-2](https://doi.org/10.1016/S0009-8981(03)00003-2)
 12. Requena JR, Chao CC, Levine RL, Stadtman ER. Glutamic and aminoadipic semialdehydes are the main carbonyl products of metal-catalyzed oxidation of proteins. *Proc Natl Acad Sci USA.* 2001; 98:69–74. <https://doi.org/10.1073/pnas.98.1.69>
 13. Kao HJ, Weng SL, Huang KY, Kaunang FJ, Hsu JB, Huang CH, Lee TY. MDD-carb: a combinatorial model for the identification of protein carbonylation sites with substrate motifs. *BMC Syst Biol.* 2017 (Suppl 7); 11:137. <https://doi.org/10.1186/s12918-017-0511-4>
 14. Suzuki YJ, Carini M, Butterfield DA. Protein carbonylation. *Antioxid Redox Signal.* 2010; 12:323–25. <https://doi.org/10.1089/ars.2009.2887>
 15. Sadowska-Bartosz I, Adamczyk-Sowa M, Galiniak S, Mucha S, Pierzchala K, Bartosz G. Oxidative modification of serum proteins in multiple sclerosis. *Neurochem Int.* 2013; 63:507–16. <https://doi.org/10.1016/j.neuint.2013.08.009>
 16. Sadowska-Bartosz I, Adamczyk-Sowa M, Gajewska A, Bartosz G. Oxidative modification of blood serum proteins in multiple sclerosis after interferon or mitoxantrone treatment. *J Neuroimmunol.* 2014; 266:67–74. <https://doi.org/10.1016/j.jneuroim.2013.11.005>
 17. Feeney MB, Schöneich C. Proteomic approaches to analyze protein tyrosine nitration. *Antioxid Redox Signal.* 2013; 19:1247–56. <https://doi.org/10.1089/ars.2012.5058>
 18. Ischiropoulos H. Biological selectivity and functional aspects of protein tyrosine nitration. *Biochem Biophys Res Commun.* 2003; 305:776–83. [https://doi.org/10.1016/S0006-291X\(03\)00814-3](https://doi.org/10.1016/S0006-291X(03)00814-3)
 19. Souza JM, Daikhin E, Yudkoff M, Raman CS, Ischiropoulos H. Factors determining the selectivity of protein tyrosine nitration. *Arch Biochem Biophys.* 1999; 371:169–78. <https://doi.org/10.1006/abbi.1999.1480>
 20. Bian K, Gao Z, Weisbrodt N, Murad F. The nature of heme/iron-induced protein tyrosine nitration. *Proc Natl Acad Sci USA.* 2003; 100:5712–17. <https://doi.org/10.1073/pnas.0931291100>
 21. Radi R. Nitric oxide, oxidants, and protein tyrosine nitration. *Proc Natl Acad Sci USA.* 2004; 101:4003–08. <https://doi.org/10.1073/pnas.0307446101>
 22. van Dalen CJ, Whitehouse MW, Winterbourn CC, Kettle AJ. Thiocyanate and chloride as competing substrates for myeloperoxidase. *Biochem J.* 1997; 327:487–92. <https://doi.org/10.1042/bj3270487>
 23. Furtmüller PG, Burner U, Obinger C. Reaction of myeloperoxidase compound I with chloride, bromide, iodide, and thiocyanate. *Biochemistry.* 1998; 37:17923–30. <https://doi.org/10.1021/bi9818772>
 24. Furtmüller PG, Zederbauer M, Jantschko W, Helm J, Bogner M, Jakopitsch C, Obinger C. Active site structure and catalytic mechanisms of human peroxidases. *Arch Biochem Biophys.* 2006; 445:199–213. <https://doi.org/10.1016/j.abb.2005.09.017>
 25. Rayner BS, Love DT, Hawkins CL. Comparative reactivity of myeloperoxidase-derived oxidants with mammalian cells. *Free Radic Biol Med.* 2014; 71:240–55. <https://doi.org/10.1016/j.freeradbiomed.2014.03.004>
 26. Li H, Cao Z, Moore DR, Jackson PL, Barnes S, Lambeth JD, Thannickal VJ, Cheng G. Microbicidal activity of vascular peroxidase 1 in human plasma via generation of hypochlorous acid. *Infect Immun.* 2012; 80:2528–37. <https://doi.org/10.1128/IAI.06337-11>
 27. Bhave G, Cummings CF, Vanacore RM, Kumagai-Cresse C, Ero-Tolliver IA, Rafi M, Kang JS, Pedchenko V, Fessler LI, Fessler JH, Hudson BG. Peroxidase forms sulfilimine chemical bonds using hypohalous acids in tissue genesis. *Nat Chem Biol.* 2012; 8:784–90. <https://doi.org/10.1038/nchembio.1038>
 28. Péterfi Z, Geiszt M. Peroxidases: novel players in tissue genesis. *Trends Biochem Sci.* 2014; 39:305–07. <https://doi.org/10.1016/j.tibs.2014.05.005>
 29. Brandes RP. Vascular peroxidase 1/peroxidase: a complex protein with a simple function? *Cardiovasc Res.* 2011;91:1-2. <https://doi.org/10.1093/cvr/cvr120>
 30. Whiteman M, Rose P, Siau JL, Cheung NS, Tan GS, Halliwell B, Armstrong JS. Hypochlorous acid-mediated mitochondrial dysfunction and apoptosis in human hepatoma HepG2 and human fetal liver cells: role of mitochondrial permeability transition. *Free Radic Biol Med.* 2005; 38:1571–84.

- <https://doi.org/10.1016/j.freeradbiomed.2005.02.030>
31. Weiss SJ, Klein R, Slivka A, Wei M. Chlorination of taurine by human neutrophils. Evidence for hypochlorous acid generation. *J Clin Invest.* 1982; 70:598–607. <https://doi.org/10.1172/JCI110652>
 32. Morris JC. The acid ionization constant of HOCl from 5 to 35°. *J Phys Chem.* 1966; 70:3798–805. <https://doi.org/10.1021/j100884a007>
 33. Arnhold J, Flemmig J. Human myeloperoxidase in innate and acquired immunity. *Arch Biochem Biophys.* 2010; 500:92–106. <https://doi.org/10.1016/j.abb.2010.04.008>
 34. Dahl JU, Gray MJ, Jakob U. Protein quality control under oxidative stress conditions. *J Mol Biol.* 2015; 427:1549–63. <https://doi.org/10.1016/j.jmb.2015.02.014>
 35. Green JN, Kettle AJ, Winterbourn CC. Protein chlorination in neutrophil phagosomes and correlation with bacterial killing. *Free Radic Biol Med.* 2014; 77:49–56. <https://doi.org/10.1016/j.freeradbiomed.2014.08.013>
 36. Pattison DI, Davies MJ. Reactions of myeloperoxidase-derived oxidants with biological substrates: gaining chemical insight into human inflammatory diseases. *Curr Med Chem.* 2006; 13:3271–90. <https://doi.org/10.2174/092986706778773095>
 37. Pattison DI, Davies MJ. Absolute rate constants for the reaction of hypochlorous acid with protein side chains and peptide bonds. *Chem Res Toxicol.* 2001; 14:1453–64. <https://doi.org/10.1021/tx0155451>
 38. Panasencko OM, Gorudko IV, Sokolov AV. Hypochlorous acid as a precursor of free radicals in living systems. *Biochemistry (Mosc).* 2013; 78:1466–89. <https://doi.org/10.1134/S0006297913130075>
 39. Dalle-Donne I, Aldini G, Carini M, Colombo R, Rossi R, Milzani A. Protein carbonylation, cellular dysfunction, and disease progression. *J Cell Mol Med.* 2006; 10:389–406. <https://doi.org/10.1111/j.1582-4934.2006.tb00407.x>
 40. Fedorova M, Bollineni RC, Hoffmann R. Protein carbonylation as a major hallmark of oxidative damage: update of analytical strategies. *Mass Spectrom Rev.* 2014; 33:79–97. <https://doi.org/10.1002/mas.21381>
 41. Aldini G, Dalle-Donne I, Facino RM, Milzani A, Carini M. Intervention strategies to inhibit protein carbonylation by lipoxidation-derived reactive carbonyls. *Med Res Rev.* 2007; 27:817–68. <https://doi.org/10.1002/med.20073>
 42. Barreiro E. Role of protein carbonylation in skeletal muscle mass loss associated with chronic conditions. *Proteomes.* 2016; 4:18. <https://doi.org/10.3390/proteomes4020018>
 43. Boden G, Homko C, Barrero CA, Stein TP, Chen X, Cheung P, Fecchio C, Koller S, Merali S. Excessive caloric intake acutely causes oxidative stress, GLUT4 carbonylation, and insulin resistance in healthy men. *Sci Translat Med.* 2015; 7:304re7. <https://doi.org/10.1126/scitranslmed.aac4765>
 44. Wong CM, Cheema AK, Zhang L, Suzuki YJ. Protein carbonylation as a novel mechanism in redox signaling. *Circ Res.* 2008; 102:310–18. <https://doi.org/10.1161/CIRCRESAHA.107.159814>
 45. Wong CM, Marcocci L, Liu L, Suzuki YJ. Cell signaling by protein carbonylation and decarbonylation. *Antioxid Redox Signal.* 2010; 12:393–404. <https://doi.org/10.1089/ars.2009.2805>
 46. Wong CM, Bansal G, Marcocci L, Suzuki YJ. Proposed role of primary protein carbonylation in cell signaling. *Redox Rep.* 2012; 17:90–94. <https://doi.org/10.1179/1351000212Y.0000000007>
 47. Oliver CN, Ahn BW, Moerman EJ, Goldstein S, Stadtman ER. Age-related changes in oxidized proteins. *J Biol Chem.* 1987; 262:5488–91.
 48. Garland D. Role of site-specific, metal-catalyzed oxidation in lens aging and cataract: a hypothesis. *Exp Eye Res.* 1990; 50:677–82. [https://doi.org/10.1016/0014-4835\(90\)90113-9](https://doi.org/10.1016/0014-4835(90)90113-9)
 49. Starke-Reed PE, Oliver CN. Protein oxidation and proteolysis during aging and oxidative stress. *Arch Biochem Biophys.* 1989; 275:559–67. [https://doi.org/10.1016/0003-9861\(89\)90402-5](https://doi.org/10.1016/0003-9861(89)90402-5)
 50. Sohal RS, Agarwal S, Dubey A, Orr WC. Protein oxidative damage is associated with life expectancy of houseflies. *Proc Natl Acad Sci USA.* 1993; 90:7255–59. <https://doi.org/10.1073/pnas.90.15.7255>
 51. Adachi H, Fujiwara Y, Ishii N. Effects of oxygen on protein carbonyl and aging in *Caenorhabditis elegans* mutants with long (*age-1*) and short (*mev-1*) life spans. *J Gerontol A Biol Sci Med Sci.* 1998; 53:B240–44. <https://doi.org/10.1093/gerona/53A.4.B240>
 52. Ahmed EK, Picot CR, Bulteau AL, Friguet B. Protein oxidative modifications and replicative senescence of WI-38 human embryonic fibroblasts. *Ann N Y Acad Sci.* 2007; 1119:88–96. <https://doi.org/10.1196/annals.1404.020>
 53. Bozaykut P, Sozen E, Kaga E, Ece A, Ozaltin E, Ek B, Ozer NK, Grune T, Bergquist J, Karademir B. The role of heat stress on the age related protein carbonylation. *J Proteomics.* 2013; 89:238–54. <https://doi.org/10.1016/j.jprot.2013.06.025>
 54. Dkhar P, Sharma R. Late-onset dietary restriction mo-

- dulates protein carbonylation and catalase in cerebral hemispheres of aged mice. *Cell Mol Neurobiol*. 2014; 34:307–13. <https://doi.org/10.1007/s10571-013-0015-8>
55. Tanase M, Urbanska AM, Zolla V, Clement CC, Huang L, Morozova K, Follo C, Goldberg M, Roda B, Reschiglian P, Santambrogio L. Role of Carbonyl Modifications on Aging-Associated Protein Aggregation. *Sci Rep*. 2016; 6:19311. <https://doi.org/10.1038/srep19311>
 56. Wu J, Luo X, Jing S, Yan LJ. Two-dimensional gel electrophoretic detection of protein carbonyls derivatized with biotin-hydrazide. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2016; 1019:128–31. <https://doi.org/10.1016/j.jchromb.2015.11.003>
 57. Feng J, Xie H, Meany DL, Thompson LV, Arriaga EA, Griffin TJ. Quantitative proteomic profiling of muscle type-dependent and age-dependent protein carbonylation in rat skeletal muscle mitochondria. *J Gerontol A Biol Sci Med Sci*. 2008; 63:1137–52. <https://doi.org/10.1093/gerona/63.11.1137>
 58. Jha R, Rizvi SI. Carbonyl formation in erythrocyte membrane proteins during aging in humans. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*. 2011; 155:39–42. <https://doi.org/10.5507/bp.2011.013>
 59. Rabek JP, Boylston WH 3rd, Papaconstantinou J. Carbonylation of ER chaperone proteins in aged mouse liver. *Biochem Biophys Res Commun*. 2003; 305:566–72. [https://doi.org/10.1016/S0006-291X\(03\)00826-X](https://doi.org/10.1016/S0006-291X(03)00826-X)
 60. Anand S, Rajashekharaiya V, Tekupalli R. Effect of age and physical activity on oxidative stress parameters in experimental rat model. *Int J Clin Exp Physiol*. 2015; 2:185–90. <https://doi.org/10.4103/2348-8093.169960>
 61. Beltran Valls MR, Wilkinson DJ, Narici MV, Smith K, Phillips BE, Caporossi D, Atherton PJ. Protein carbonylation and heat shock proteins in human skeletal muscle: relationships to age and sarcopenia. *J Gerontol A Biol Sci Med Sci*. 2015; 70:174–81. <https://doi.org/10.1093/gerona/glu007>
 62. Wang Z, Wang Y, Liu H, Che Y, Xu Y, e L. Age-related variations of protein carbonyls in human saliva and plasma: is saliva protein carbonyls an alternative biomarker of aging? *Age (Dordr)*. 2015; 37:9781. <https://doi.org/10.1007/s11357-015-9781-1>
 63. Tohma H, El-Shafey AF, Croft K, Shavlakadze T, Grounds MD, Arthur PG. Protein thiol oxidation does not change in skeletal muscles of aging female mice. *Biogerontology*. 2014; 15:87–98. <https://doi.org/10.1007/s10522-013-9483-y>
 64. Rowiński R, Kozakiewicz M, Kędziora-Kornatowska K, Hübner-Woźniak E, Kędziora J. Markers of oxidative stress and erythrocyte antioxidant enzyme activity in older men and women with differing physical activity. *Exp Gerontol*. 2013; 48:1141–46. <https://doi.org/10.1016/j.exger.2013.07.010>
 65. Lourenço dos Santos S, Baraibar MA, Lundberg S, Eeg-Olofsson O, Larsson L, Friguet B. Oxidative proteome alterations during skeletal muscle ageing. *Redox Biol*. 2015; 5:267–74. <https://doi.org/10.1016/j.redox.2015.05.006>
 66. Barreiro E, Coronell C, Laviña B, Ramírez-Sarmiento A, Orozco-Levi M, Gea J, and PENAM Project. Aging, sex differences, and oxidative stress in human respiratory and limb muscles. *Free Radic Biol Med*. 2006; 41:797–809. <https://doi.org/10.1016/j.freeradbiomed.2006.05.027>
 67. Stadtman ER. Metal ion-catalyzed oxidation of proteins: biochemical mechanism and biological consequences. *Free Radic Biol Med*. 1990; 9:315–25. [https://doi.org/10.1016/0891-5849\(90\)90006-5](https://doi.org/10.1016/0891-5849(90)90006-5)
 68. Sohal RS, Agarwal A, Agarwal S, Orr WC. Simultaneous overexpression of copper- and zinc-containing superoxide dismutase and catalase retards age-related oxidative damage and increases metabolic potential in *Drosophila melanogaster*. *J Biol Chem*. 1995; 270:15671–74. <https://doi.org/10.1074/jbc.270.26.15671>
 69. Dubey A, Forster MJ, Lal H, Sohal RS. Effect of age and caloric intake on protein oxidation in different brain regions and on behavioral functions of the mouse. *Arch Biochem Biophys*. 1996; 333:189–97. <https://doi.org/10.1006/abbi.1996.0380>
 70. Forster MJ, Sohal BH, Sohal RS. Reversible effects of long-term caloric restriction on protein oxidative damage. *J Gerontol A Biol Sci Med Sci*. 2000; 55:B522–29. <https://doi.org/10.1093/gerona/55.11.B522>
 71. Levine RL. Carbonyl modified proteins in cellular regulation, aging, and disease. *Free Radic Biol Med*. 2002; 32:790–96. [https://doi.org/10.1016/S0891-5849\(02\)00765-7](https://doi.org/10.1016/S0891-5849(02)00765-7)
 72. Sohal RS. Role of oxidative stress and protein oxidation in the aging process. *Free Radic Biol Med*. 2002; 33:37–44. [https://doi.org/10.1016/S0891-5849\(02\)00856-0](https://doi.org/10.1016/S0891-5849(02)00856-0)
 73. Choi J, Rees HD, Weintraub ST, Levey AI, Chin LS, Li L. Oxidative modifications and aggregation of Cu,Zn-superoxide dismutase associated with Alzheimer and Parkinson diseases. *J Biol Chem*. 2005; 280:11648–55. <https://doi.org/10.1074/jbc.M414327200>

74. Das N, Levine RL, Orr WC, Sohal RS. Selectivity of protein oxidative damage during aging in *Drosophila melanogaster*. *Biochem J*. 2001; 360:209–16. <https://doi.org/10.1042/bj3600209>
75. Jana CK, Das N, Sohal RS. Specificity of age-related carbonylation of plasma proteins in the mouse and rat. *Arch Biochem Biophys*. 2002; 397:433–39. <https://doi.org/10.1006/abbi.2001.2690>
76. Maisonneuve E, Ducret A, Khoueiry P, Lignon S, Longhi S, Talla E, Dukan S. Rules governing selective protein carbonylation. *PLoS One*. 2009; 4:e7269. <https://doi.org/10.1371/journal.pone.0007269>
77. Rommer PS, Greilberger J, Salhofer-Polanyi S, Auff E, Leutmezer F, Herwig R. Elevated levels of carbonyl proteins in cerebrospinal fluid of patients with neurodegenerative diseases. *Tohoku J Exp Med*. 2014; 234:313–17. <https://doi.org/10.1620/tjem.234.313>
78. Adamczyk-Sowa M, Bieszczad-Bedrejcuk E, Galiniak S, Rozmiłowska I, Czyżewski D, Bartosz G, Sadowska-Bartosz I. Oxidative modifications of blood serum proteins in myasthenia gravis. *J Neuroimmunol*. 2017; 305:145–53. <https://doi.org/10.1016/j.jneuroim.2017.01.019>
79. Bizzozero OA, DeJesus G, Callahan K, Pastuszyn A. Elevated protein carbonylation in the brain white matter and gray matter of patients with multiple sclerosis. *J Neurosci Res*. 2005; 81:687–95. <https://doi.org/10.1002/jnr.20587>
80. Hensley K, Hall N, Subramaniam R, Cole P, Harris M, Aksenov M, Aksenova M, Gabbita SP, Wu JF, Carney JM, Lovell M, Markesbery WR, Butterfield DA. Brain regional correspondence between Alzheimer's disease histopathology and biomarkers of protein oxidation. *J Neurochem*. 1995; 65:2146–56. <https://doi.org/10.1046/j.1471-159.1995.65052146.x>
81. Butterfield DA, Perluigi M, Reed T, Muharib T, Hughes CP, Robinson RA, Sultana R. Redox proteomics in selected neurodegenerative disorders: from its infancy to future applications. *Antioxid Redox Signal*. 2012; 17:1610–55. <https://doi.org/10.1089/ars.2011.4109>
82. Sultana R, Boyd-Kimball D, Poon HF, Cai J, Pierce WM, Klein JB, Merchant M, Markesbery WR, Butterfield DA. Redox proteomics identification of oxidized proteins in Alzheimer's disease hippocampus and cerebellum: an approach to understand pathological and biochemical alterations in AD. *Neurobiol Aging*. 2006; 27:1564–76. <https://doi.org/10.1016/j.neurobiolaging.2005.09.021>
83. Butterfield DA, Sultana R. Redox proteomics identification of oxidatively modified brain proteins in Alzheimer's disease and mild cognitive impairment: insights into the progression of this dementing disorder. *J Alzheimers Dis*. 2007; 12:61–72. <https://doi.org/10.3233/JAD-2007-12107>
84. Aluise CD, Robinson RA, Cai J, Pierce WM, Markesbery WR, Butterfield DA. Redox proteomics analysis of brains from subjects with amnesic mild cognitive impairment compared to brains from subjects with preclinical Alzheimer's disease: insights into memory loss in MCI. *J Alzheimers Dis*. 2011; 23:257–69. <https://doi.org/10.3233/JAD-2010-10108>
85. Zabel M, Nackenoff A, Kirsch WM, Harrison FE, Perry G, Schrag M. Markers of oxidative damage to lipids, nucleic acids and proteins and antioxidant enzymes activities in Alzheimer's disease brain: A meta-analysis in human pathological specimens. *Free Radic Biol Med*. 2018; 115:351–60. <https://doi.org/10.1016/j.freeradbiomed.2017.12.016>
86. Shen L, Chen Y, Yang A, Chen C, Liao L, Li S, Ying M, Tian J, Liu Q, Ni J. Redox Proteomic Profiling of Specifically Carbonylated Proteins in the Serum of Triple Transgenic Alzheimer's Disease Mice. *Int J Mol Sci*. 2016; 17:469. <https://doi.org/10.3390/ijms17040469>
87. Sorolla MA, Rodríguez-Colman MJ, Tamarit J, Ortega Z, Lucas JJ, Ferrer I, Ros J, Cabiscol E. Protein oxidation in Huntington disease affects energy production and vitamin B6 metabolism. *Free Radic Biol Med*. 2010; 49:612–21. <https://doi.org/10.1016/j.freeradbiomed.2010.05.016>
88. Butterfield DA, Gu L, Di Domenico F, Robinson RA. Mass spectrometry and redox proteomics: applications in disease. *Mass Spectrom Rev*. 2014; 33:277–301. <https://doi.org/10.1002/mas.21374>
89. Liu Q, Raina AK, Smith MA, Sayre LM, Perry G. Hydroxynonenal, toxic carbonyls, and Alzheimer disease. *Mol Aspects Med*. 2003; 24:305–13. [https://doi.org/10.1016/S0098-2997\(03\)00025-6](https://doi.org/10.1016/S0098-2997(03)00025-6)
90. McGrath LT, McGleenon BM, Brennan S, McColl D, McLroy S, Passmore AP. Increased oxidative stress in Alzheimer's disease as assessed with 4-hydroxynonenal but not malondialdehyde. *QJM*. 2001; 94:485–90. <https://doi.org/10.1093/qjmed/94.9.485>
91. Selley ML. (E)-4-hydroxy-2-nonenal may be involved in the pathogenesis of Parkinson's disease. *Free Radic Biol Med*. 1998; 25:169–74. [https://doi.org/10.1016/S0891-5849\(98\)00021-5](https://doi.org/10.1016/S0891-5849(98)00021-5)
92. Dei R, Takeda A, Niwa H, Li M, Nakagomi Y, Watanabe M, Inagaki T, Washimi Y, Yasuda Y, Horie K, Miyata T, Sobue G. Lipid peroxidation and advanced glycation end products in the brain in normal aging and in

- Alzheimer's disease. *Acta Neuropathol.* 2002; 104:113–22. <https://doi.org/10.1007/s00401-002-0523-y>
93. Simpson EP, Henry YK, Henkel JS, Smith RG, Appel SH. Increased lipid peroxidation in sera of ALS patients: a potential biomarker of disease burden. *Neurology.* 2004; 62:1758–65. <https://doi.org/10.1212/WNL.62.10.1758>
94. Wehr NB, Levine RL. Quantification of protein carbonylation. *Methods Mol Biol.* 2013; 965:265–81. https://doi.org/10.1007/978-1-62703-239-1_18
95. Mesquita CS, Oliveira R, Bento F, Geraldo D, Rodrigues JV, Marcos JC. Simplified 2,4-dinitrophenylhydrazine spectrophotometric assay for quantification of carbonyls in oxidized proteins. *Anal Biochem.* 2014; 458:69–71. <https://doi.org/10.1016/j.ab.2014.04.034>
96. Wehr NB, Levine RL. Quantitation of protein carbonylation by dot blot. *Anal Biochem.* 2012; 423:241–45. <https://doi.org/10.1016/j.ab.2012.01.031>
97. Pazos M, da Rocha AP, Roepstorff P, Rogowska-Wrzesinska A. Fish proteins as targets of ferrous-catalyzed oxidation: identification of protein carbonyls by fluorescent labeling on two-dimensional gels and MALDI-TOF/TOF mass spectrometry. *J Agric Food Chem.* 2011; 59:7962–77. <https://doi.org/10.1021/jf201080t>
98. Tamarit J, de Hoogh A, Obis E, Alsina D, Cabiscol E, Ros J. Analysis of oxidative stress-induced protein carbonylation using fluorescent hydrazides. *J Proteomics.* 2012; 75:3778–88. <https://doi.org/10.1016/j.jprot.2012.04.046>
99. Pyr Dit Ruys S, Bonzom JM, Frelon S. Benchmarking of protein carbonylation analysis in *Caenorhabditis elegans*: specific considerations and general advice. *Free Radic Biol Med.* 2016; 99:364–73. <https://doi.org/10.1016/j.freeradbiomed.2016.08.014>
100. Mukherjee K, Chio TI, Sackett DL, Bane SL. Detection of oxidative stress-induced carbonylation in live mammalian cells. *Free Radic Biol Med.* 2015; 84:11–21. <https://doi.org/10.1016/j.freeradbiomed.2015.03.011>
101. Artemenko K, Mi J, Bergquist J. Mass-spectrometry-based characterization of oxidations in proteins. *Free Radic Res.* 2015; 49:477–93. <https://doi.org/10.3109/10715762.2015.1023795>
102. Havelund JF, Wojdyla K, Davies MJ, Jensen ON, Møller IM, Rogowska-Wrzesinska A. A biotin enrichment strategy identifies novel carbonylated amino acids in proteins from human plasma. *J Proteomics.* 2017; 156:40–51. <https://doi.org/10.1016/j.jprot.2016.12.019>
103. Radi R. Protein tyrosine nitration: biochemical mechanisms and structural basis of functional effects. *Acc Chem Res.* 2013; 46:550–59. <https://doi.org/10.1021/ar300234c>
104. Ischiropoulos H. Protein tyrosine nitration--an update. *Arch Biochem Biophys.* 2009; 484:117–21. <https://doi.org/10.1016/j.abb.2008.10.034>
105. Chakravarti B, Chakravarti DN. Protein Tyrosine Nitration: role in Aging. *Curr Aging Sci.* 2017; 10:246–62. <https://doi.org/10.2174/1874609810666170315112634>
106. Yeo WS, Kim YJ, Kabir MH, Kang JW, Ahsan-Ul-Bari M, Kim KP. Mass spectrometric analysis of protein tyrosine nitration in aging and neurodegenerative diseases. *Mass Spectrom Rev.* 2015; 34:166–83. <https://doi.org/10.1002/mas.21429>
107. Masri F. Role of nitric oxide and its metabolites as potential markers in lung cancer. *Ann Thorac Med.* 2010; 5:123–27. <https://doi.org/10.4103/1817-1737.65036>
108. Bradley SA, Steinert JR. Nitric Oxide-Mediated Posttranslational Modifications: impacts at the Synapse. *Oxid Med Cell Longev.* 2016; 2016:5681036. <https://doi.org/10.1155/2016/5681036>
109. Kummer MP, Hermes M, Delekarte A, Hammerschmidt T, Kumar S, Terwel D, Walter J, Pape HC, König S, Roeber S, Jessen F, Klockgether T, Korte M, Heneka MT. Nitration of tyrosine 10 critically enhances amyloid β aggregation and plaque formation. *Neuron.* 2011; 71:833–44. <https://doi.org/10.1016/j.neuron.2011.07.001>
110. Calabrese V, Boyd-Kimball D, Scapagnini G, Butterfield DA. Nitric oxide and cellular stress response in brain aging and neurodegenerative disorders: the role of vitagenes. *In Vivo.* 2004; 18:245–67.
111. Aquilano K, Baldelli S, Cardaci S, Rotilio G, Ciriolo MR. Nitric oxide is the primary mediator of cytotoxicity induced by GSH depletion in neuronal cells. *J Cell Sci.* 2011; 124:1043–54. <https://doi.org/10.1242/jcs.077149>
112. Solano RM, Casarejos MJ, Menéndez-Cuervo J, Rodríguez-Navarro JA, García de Yébenes J, Mena MA. Glial dysfunction in parkin null mice: effects of aging. *J Neurosci.* 2008; 28:598–611. <https://doi.org/10.1523/JNEUROSCI.4609-07.2008>
113. Ballatori N, Krance SM, Notenboom S, Shi S, Tieu K, Hammond CL. Glutathione dysregulation and the etiology and progression of human diseases. *Biol Chem.* 2009; 390:191–214.

<https://doi.org/10.1515/BC.2009.033>

114. Mallozzi C, D'Amore C, Camerini S, Macchia G, Crescenzi M, Petrucci TC, Di Stasi AM. Phosphorylation and nitration of tyrosine residues affect functional properties of Synaptophysin and Dynamin I, two proteins involved in exo-endocytosis of synaptic vesicles. *Biochim Biophys Acta*. 2013; 1833:110–21. <https://doi.org/10.1016/j.bbamcr.2012.10.022>
115. Di Stasi AM, Mallozzi C, Macchia G, Maura G, Petrucci TC, Minetti M. Peroxynitrite affects exocytosis and SNARE complex formation and induces tyrosine nitration of synaptic proteins. *J Neurochem*. 2002; 82:420–29. <https://doi.org/10.1046/j.1471-4159.2002.00980.x>
116. Mallozzi C, Ceccarini M, Camerini S, Macchia G, Crescenzi M, Petrucci TC, Di Stasi AM. Peroxynitrite induces tyrosine residue modifications in synaptophysin C-terminal domain, affecting its interaction with src. *J Neurochem*. 2009; 111:859–69. <https://doi.org/10.1111/j.1471-4159.2009.06378.x>
117. Vrljic M, Strop P, Hill RC, Hansen KC, Chu S, Brunger AT. Post-translational modifications and lipid binding profile of insect cell-expressed full-length mammalian synaptotagmin 1. *Biochemistry*. 2011; 50:9998–10012. <https://doi.org/10.1021/bi200998y>
118. Tran MH, Yamada K, Olariu A, Mizuno M, Ren XH, Nabeshima T. Amyloid beta-peptide induces nitric oxide production in rat hippocampus: association with cholinergic dysfunction and amelioration by inducible nitric oxide synthase inhibitors. *FASEB J*. 2001; 15:1407–09. <https://doi.org/10.1096/fj.00-0719fje>
119. Tran MH, Yamada K, Nakajima A, Mizuno M, He J, Kamei H, Nabeshima T. Tyrosine nitration of a synaptic protein synaptophysin contributes to amyloid beta-peptide-induced cholinergic dysfunction. *Mol Psychiatry*. 2003; 8:407–12. <https://doi.org/10.1038/sj.mp.4001240>
120. Butterfield DA, Reed TT, Perluigi M, De Marco C, Coccia R, Keller JN, Markesbery WR, Sultana R. Elevated levels of 3-nitrotyrosine in brain from subjects with amnesic mild cognitive impairment: implications for the role of nitration in the progression of Alzheimer's disease. *Brain Res*. 2007; 1148:243–48. <https://doi.org/10.1016/j.brainres.2007.02.084>
121. Halliwell B. What nitrates tyrosine? Is nitrotyrosine specific as a biomarker of peroxynitrite formation in vivo? *FEBS Lett*. 1997; 411:157–60. [https://doi.org/10.1016/S0014-5793\(97\)00469-9](https://doi.org/10.1016/S0014-5793(97)00469-9)
122. Horiguchi T, Uryu K, Giasson BI, Ischiropoulos H, Lightfoot R, Bellmann C, Richter-Landsberg C, Lee VM, Trojanowski JQ. Nitration of tau protein is linked to neurodegeneration in tauopathies. *Am J Pathol*. 2003; 163:1021–31. [https://doi.org/10.1016/S0002-9440\(10\)63462-1](https://doi.org/10.1016/S0002-9440(10)63462-1)
123. Reynolds MR, Reyes JF, Fu Y, Bigio EH, Guillozet-Bongaarts AL, Berry RW, Binder LI. Tau nitration occurs at tyrosine 29 in the fibrillar lesions of Alzheimer's disease and other tauopathies. *J Neurosci*. 2006; 26:10636–45. <https://doi.org/10.1523/JNEUROSCI.2143-06.2006>
124. Reyes JF, Fu Y, Vana L, Kanaan NM, Binder LI. Tyrosine nitration within the proline-rich region of Tau in Alzheimer's disease. *Am J Pathol*. 2011; 178:2275–85. <https://doi.org/10.1016/j.ajpath.2011.01.030>
125. Adav SS, Sze SK. Insight of brain degenerative protein modifications in the pathology of neurodegeneration and dementia by proteomic profiling. *Mol Brain*. 2016; 9:92. <https://doi.org/10.1186/s13041-016-0272-9>
126. Thomas SN, Yang AJ. Mass spectrometry analysis of lysine posttranslational modifications of tau protein from Alzheimer's disease brain. *Methods Mol Biol*. 2017; 1523:161–77. https://doi.org/10.1007/978-1-4939-6598-4_10
127. Yuan C, Yi L, Yang Z, Deng Q, Huang Y, Li H, Gao Z. Amyloid beta-heme peroxidase promoted protein nitrotyrosination: relevance to widespread protein nitration in Alzheimer's disease. *J Biol Inorg Chem*. 2012; 17:197–207. <https://doi.org/10.1007/s00775-011-0842-3>
128. Guix FX, Ill-Raga G, Bravo R, Nakaya T, de Fabritiis G, Coma M, Miscione GP, Villà-Freixa J, Suzuki T, Fernández-Busquets X, Valverde MA, de Strooper B, Muñoz FJ. Amyloid-dependent triosephosphate isomerase nitrotyrosination induces glycation and tau fibrillation. *Brain*. 2009; 132:1335–45. <https://doi.org/10.1093/brain/awp023>
129. Castegna A, Thongboonkerd V, Klein JB, Lynn B, Markesbery WR, Butterfield DA. Proteomic identification of nitrated proteins in Alzheimer's disease brain. *J Neurochem*. 2003; 85:1394–401. <https://doi.org/10.1046/j.1471-4159.2003.01786.x>
130. Sultana R, Poon HF, Cai J, Pierce WM, Merchant M, Klein JB, Markesbery WR, Butterfield DA. Identification of nitrated proteins in Alzheimer's disease brain using a redox proteomics approach. *Neurobiol Dis*. 2006; 22:76–87. <https://doi.org/10.1016/j.nbd.2005.10.004>
131. Reed TT, Pierce WM Jr, Turner DM, Markesbery WR, Butterfield DA. Proteomic identification of nitrated brain proteins in early Alzheimer's disease inferior parietal lobule. *J Cell Mol Med*. 2009; 13:2019–29.

- <https://doi.org/10.1111/j.1582-4934.2008.00478.x>
132. McCormack AL, Mak SK, Di Monte DA. Increased α -synuclein phosphorylation and nitration in the aging primate substantia nigra. *Cell Death Dis.* 2012; 3:e315. <https://doi.org/10.1038/cddis.2012.50>
133. Schildknecht S, Gerding HR, Karreman C, Drescher M, Lashuel HA, Outeiro TF, Di Monte DA, Leist M. Oxidative and nitrative alpha-synuclein modifications and proteostatic stress: implications for disease mechanisms and interventions in synucleinopathies. *J Neurochem.* 2013; 125:491–511. <https://doi.org/10.1111/jnc.12226>
134. Marques O, Outeiro TF. Alpha-synuclein: from secretion to dysfunction and death. *Cell Death Dis.* 2012; 3:e350. <https://doi.org/10.1038/cddis.2012.94>
135. Oliveira LM, Falomir-Lockhart LJ, Botelho MG, Lin KH, Wales P, Koch JC, Gerhardt E, Taschenberger H, Outeiro TF, Lingor P, Schüle B, Arndt-Jovin DJ, Jovin TM. Elevated α -synuclein caused by SNCA gene triplication impairs neuronal differentiation and maturation in Parkinson's patient-derived induced pluripotent stem cells. *Cell Death Dis.* 2015; 6:e1994. <https://doi.org/10.1038/cddis.2015.318>
136. Burai R, Ait-Bouziad N, Chiki A, Lashuel HA. Elucidating the Role of Site-Specific Nitration of α -Synuclein in the Pathogenesis of Parkinson's Disease via Protein Semisynthesis and Mutagenesis. *J Am Chem Soc.* 2015; 137:5041–52. <https://doi.org/10.1021/ja5131726>
137. Benner EJ, Banerjee R, Reynolds AD, Sherman S, Pisarev VM, Tsiperson V, Nemachek C, Ciborowski P, Przedborski S, Mosley RL, Gendelman HE. Nitrated alpha-synuclein immunity accelerates degeneration of nigral dopaminergic neurons. *PLoS One.* 2008; 3:e1376. <https://doi.org/10.1371/journal.pone.0001376>
138. Prigione A, Piazza F, Brighina L, Begni B, Galbussera A, Difrancesco JC, Andreoni S, Piolti R, Ferrarese C. Alpha-synuclein nitration and autophagy response are induced in peripheral blood cells from patients with Parkinson disease. *Neurosci Lett.* 2010; 477:6–10. <https://doi.org/10.1016/j.neulet.2010.04.022>
139. Kleinknecht A, Popova B, Lázaro DF, Pinho R, Valerius O, Outeiro TF, Braus GH. C-Terminal Tyrosine Residue Modifications Modulate the Protective Phosphorylation of Serine 129 of α -Synuclein in a Yeast Model of Parkinson's Disease. *PLoS Genet.* 2016; 12:e1006098. <https://doi.org/10.1371/journal.pgen.1006098>
140. Kim YJ, Bari AU, Kim KP. Mass Spectrometric Analysis of Protein Tyrosine Nitration in Aging and Relevant Diseases. *Bio Design.* 2013; 1:26–31.
141. Peluffo G, Radi R. Biochemistry of protein tyrosine nitration in cardiovascular pathology. *Cardiovasc Res.* 2007; 75:291–302. <https://doi.org/10.1016/j.cardiores.2007.04.024>
142. Thomas SN, Yang AJ. Mass Spectrometry Analysis of Lysine Posttranslational Modifications of Tau Protein from Alzheimer's Disease Brain. *Methods Mol Biol.* 2017; 1523:161–77. https://doi.org/10.1007/978-1-4939-6598-4_10
143. Ray RS, Katyal A. Myeloperoxidase: bridging the gap in neurodegeneration. *Neurosci Biobehav Rev.* 2016; 68:611–20. <https://doi.org/10.1016/j.neubiorev.2016.06.031>
144. Jeitner TM, Kalogiannis M, Krasnikov BF, Gomolin I, Peltier MR, Moran GR. Linking inflammation and Parkinson disease: hypochlorous acid generates Parkinsonian poisons. *Toxicol Sci.* 2016; 151:388–402. <https://doi.org/10.1093/toxsci/kfw052>
145. Lefkowitz DL, Lefkowitz SS. Microglia and myeloperoxidase: a deadly partnership in neurodegenerative disease. *Free Radic Biol Med.* 2008; 45:726–31. <https://doi.org/10.1016/j.freeradbiomed.2008.05.021>
146. Choi DK, Pennathur S, Perier C, Tieu K, Teismann P, Wu DC, Jackson-Lewis V, Vila M, Vonsattel JP, Heinecke JW, Przedborski S. Ablation of the inflammatory enzyme myeloperoxidase mitigates features of Parkinson's disease in mice. *J Neurosci.* 2005; 25:6594–600. <https://doi.org/10.1523/JNEUROSCI.0970-05.2005>
147. Reynolds WF, Rhee J, Maciejewski D, Paladino T, Sieburg H, Maki RA, Masliah E. Myeloperoxidase polymorphism is associated with gender specific risk for Alzheimer's disease. *Exp Neurol.* 1999; 155:31–41. <https://doi.org/10.1006/exnr.1998.6977>
148. Nagra RM, Becher B, Tourtellotte WW, Antel JP, Gold D, Paladino T, Smith RA, Nelson JR, Reynolds WF. Immunohistochemical and genetic evidence of myeloperoxidase involvement in multiple sclerosis. *J Neuroimmunol.* 1997; 78:97–107. [https://doi.org/10.1016/S0165-5728\(97\)00089-1](https://doi.org/10.1016/S0165-5728(97)00089-1)
149. Boven LA, Middel J, Verhoef J, De Groot CJ, Nottet HS. Monocyte infiltration is highly associated with loss of the tight junction protein zonula occludens in HIV-1-associated dementia. *Neuropathol Appl Neurobiol.* 2000; 26:356–60. <https://doi.org/10.1046/j.1365-2990.2000.00255.x>
150. Green PS, Mendez AJ, Jacob JS, Crowley JR, Growdon W, Hyman BT, Heinecke JW. Neuronal expression of myeloperoxidase is increased in Alzheimer's disease. *J Neurochem.* 2004; 90:724–33. <https://doi.org/10.1111/j.1471-4159.2004.02527.x>

151. Chang ML, Klaidman LK Jr, Adams JD Jr. The effects of oxidative stress on in vivo brain GSH turnover in young and mature mice. *Mol Chem Neuropathol*. 1997; 30:187–97. <https://doi.org/10.1007/BF02815097>
152. Eto K, Asada T, Arima K, Makifuchi T, Kimura H. Brain hydrogen sulfide is severely decreased in Alzheimer's disease. *Biochem Biophys Res Commun*. 2002; 293:1485–88. [https://doi.org/10.1016/S0006-291X\(02\)00422-9](https://doi.org/10.1016/S0006-291X(02)00422-9)
153. Krasowska A, Konat GW. Vulnerability of brain tissue to inflammatory oxidant, hypochlorous acid. *Brain Res*. 2004; 997:176–84. <https://doi.org/10.1016/j.brainres.2003.09.080>
154. Yap YW, Whiteman M, Bay BH, Li Y, Sheu FS, Qi RZ, Tan CH, Cheung NS. Hypochlorous acid induces apoptosis of cultured cortical neurons through activation of calpains and rupture of lysosomes. *J Neurochem*. 2006; 98:1597–609. <https://doi.org/10.1111/j.1471-4159.2006.03996.x>
155. Yap YW, Chen MJ, Choy MS, Peng ZF, Whiteman M, Manikandan J, Melendez AJ, Cheung NS. Temporal transcriptomic profiling reveals cellular targets that govern survival in HOCl-mediated neuronal apoptosis. *Life Sci*. 2010; 87:457–67. <https://doi.org/10.1016/j.lfs.2010.08.011>
156. Casciaro M, Di Salvo E, Pace E, Ventura-Spagnolo E, Navarra M, Gangemi S. Chlorinative stress in age-related diseases: a literature review. *Immun Ageing*. 2017; 14:21. <https://doi.org/10.1186/s12979-017-0104-5>
157. Pizzi A, Pigliacelli C, Gori A, Nonappa, Ikkala O, Demitri N, Terraneo G, Castelletto V, Hamley IW, Baldelli Bombelli F, Metrangolo P. Halogenation dictates the architecture of amyloid peptide nanostructures. *Nanoscale*. 2017; 9:9805–10. <https://doi.org/10.1039/C7NR03263C>
158. Witko-Sarsat V, Friedlander M, Nguyen Khoa T, Capeillère-Blandin C, Nguyen AT, Canteloup S, Dayer JM, Jungers P, Drüeke T, Descamps-Latscha B. Advanced oxidation protein products as novel mediators of inflammation and monocyte activation in chronic renal failure. *J Immunol*. 1998; 161:2524–32.
159. Cao W, Hou FF, Nie J. AOPPs and the progression of kidney disease. *Kidney Int Suppl* (2011). 2014; 4:102–06. <https://doi.org/10.1038/kisup.2014.19>
160. Witko-Sarsat V, Friedlander M, Capeillère-Blandin C, Nguyen-Khoa T, Nguyen AT, Zingraff J, Jungers P, Descamps-Latscha B. Advanced oxidation protein products as a novel marker of oxidative stress in uremia. *Kidney Int*. 1996; 49:1304–13. <https://doi.org/10.1038/ki.1996.186>
161. Atukeren P, Cengiz M, Yavuzer H, Gelisgen R, Altunoglu E, Oner S, Erdenen F, Yucaekin D, Derici H, Cakatay U, Uzun H. The efficacy of donepezil administration on acetylcholinesterase activity and altered redox homeostasis in Alzheimer's disease. *Biomed Pharmacother*. 2017; 90:786–95. <https://doi.org/10.1016/j.biopha.2017.03.101>
162. Boll KM, Noto C, Bonifácio KL, Bortolaschi CC, Gadelha A, Bressan RA, Barbosa DS, Maes M, Moreira EG. Oxidative and nitrosative stress biomarkers in chronic schizophrenia. *Psychiatry Res*. 2017; 253:43–48. <https://doi.org/10.1016/j.psychres.2017.03.038>
163. Medeiros MS, Schumacher-Schuh A, Cardoso AM, Bochi GV, Baldissarelli J, Kegler A, Santana D, Chaves CM, Schetinger MR, Moresco RN, Rieder CR, Figuera MR. Iron and Oxidative Stress in Parkinson's Disease: An Observational Study of Injury Biomarkers. *PLoS One*. 2016; 11:e0146129. <https://doi.org/10.1371/journal.pone.0146129>
164. Cakir T, Goktas B, Mutlu MF, Mutlu I, Bilgihan A, Erdem M, Erdem A. Advanced oxidation protein products and malondialdehyde - the new biological markers of oxidative stress - are elevated in post-menopausal women. *Ginekol Pol*. 2016; 87:321–25. <https://doi.org/10.5603/GP.2016.0001>
165. Allanore Y, Borderie D, Lemaréchal H, Ekindjian OG, Kahan A. Nifedipine decreases sVCAM-1 concentrations and oxidative stress in systemic sclerosis but does not affect the concentrations of vascular endothelial growth factor or its soluble receptor 1. *Arthritis Res Ther*. 2004; 6:R309–14. <https://doi.org/10.1186/ar1183>
166. Baskol G, Gumus K, Oner A, Arda H, Karakucuk S. The role of advanced oxidation protein products and total thiols in diabetic retinopathy. *Eur J Ophthalmol*. 2008; 18:792–98. <https://doi.org/10.1177/112067210801800521>
167. Hou G, Lu H, Chen M, Yao H, Zhao H. Oxidative stress participates in age-related changes in rat lumbar intervertebral discs. *Arch Gerontol Geriatr*. 2014; 59:665–69. <https://doi.org/10.1016/j.archger.2014.07.002>
168. Zhang YB, Zhong ZM, Hou G, Jiang H, Chen JT. Involvement of oxidative stress in age-related bone loss. *J Surg Res*. 2011; 169:e37–42. <https://doi.org/10.1016/j.jss.2011.02.033>
169. Cao W, Xu J, Zhou ZM, Wang GB, Hou FF, Nie J. Advanced oxidation protein products activate intrarenal renin-angiotensin system via a CD36-mediated, redox-dependent pathway. *Antioxid Redox Signal*. 2013; 18:19–35.

<https://doi.org/10.1089/ars.2012.4603>

170. Taylor EL, Armstrong KR, Perrett D, Hattersley AT, Winyard PG. Optimisation of an Advanced Oxidation Protein Products Assay: Its Application to Studies of Oxidative Stress in Diabetes Mellitus. *Oxid Med Cell Longev*. 2015; 2015:496271. <https://doi.org/10.1155/2015/496271>
171. Selmeçi L, Seres L, Soós P, Székely M, Acsády G. Kinetic assay for the determination of the oxidative stress biomarker, advanced oxidation protein products (AOPP) in the human blood plasma. *Acta Physiol Hung*. 2008; 95:209–18. <https://doi.org/10.1556/APhysiol.95.2008.2.5>
172. Oguz O, Inal BB, Emre T, Ozcan O, Altunoglu E, Oguz G, Topkaya C, Guvenen G. Is automated kinetic measurement superior to end-point for advanced oxidation protein product? *Clin Lab*. 2014; 60:925–30. <https://doi.org/10.7754/Clin.Lab.2013.130727>
173. Spickett CM. The lipid peroxidation product 4-hydroxy-2-nonenal: advances in chemistry and analysis. *Redox Biol*. 2013; 1:145–52. <https://doi.org/10.1016/j.redox.2013.01.007>
174. Zhang H, Forman HJ. 4-hydroxynonenal-mediated signaling and aging. *Free Radic Biol Med*. 2017; 111:219–25. <https://doi.org/10.1016/j.freeradbiomed.2016.11.032>
175. Esterbauer H, Benedetti A, Lang J, Fulceri R, Fauler G, Comporti M. Studies on the mechanism of formation of 4-hydroxynonenal during microsomal lipid peroxidation. *Biochim Biophys Acta*. 1986; 876:154–66. [https://doi.org/10.1016/0005-2760\(86\)90329-2](https://doi.org/10.1016/0005-2760(86)90329-2)
176. Dalleau S, Baradat M, Guéraud F, Huc L. Cell death and diseases related to oxidative stress: 4-hydroxynonenal (HNE) in the balance. *Cell Death Differ*. 2013; 20:1615–30. <https://doi.org/10.1038/cdd.2013.138>
177. Csala M, Kardon T, Legeza B, Lizák B, Mandl J, Margittai É, Puskás F, Száraz P, Szelényi P, Bánhegyi G. On the role of 4-hydroxynonenal in health and disease. *Biochim Biophys Acta*. 2015; 1852:826–38. <https://doi.org/10.1016/j.bbadis.2015.01.015>
178. Esterbauer H, Benedetti A, Lang J, Fulceri R, Fauler G, Comporti M. Studies on the mechanism of formation of 4-hydroxynonenal during microsomal lipid peroxidation. *Biochim Biophys Acta*. 1986; 876:154–66. [https://doi.org/10.1016/0005-2760\(86\)90329-2](https://doi.org/10.1016/0005-2760(86)90329-2)
179. Zarkovic N. 4-hydroxynonenal as a bioactive marker of pathophysiological processes. *Mol Aspects Med*. 2003; 24:281–91. [https://doi.org/10.1016/S0098-2997\(03\)00023-2](https://doi.org/10.1016/S0098-2997(03)00023-2)
180. Zarkovic K, Jakovcevic A, Zarkovic N. Contribution of the HNE-immunohistochemistry to modern pathological concepts of major human diseases. *Free Radic Biol Med*. 2017; 111:110–26. <https://doi.org/10.1016/j.freeradbiomed.2016.12.009>
181. Robino G, Parola M, Marra F, Caligiuri A, De Franco RM, Zamara E, Bellomo G, Gentilini P, Pinzani M, Dianzani MU. Interaction between 4-hydroxy-2,3-alkenals and the platelet-derived growth factor-beta receptor. Reduced tyrosine phosphorylation and downstream signaling in hepatic stellate cells. *J Biol Chem*. 2000; 275:40561–67. <https://doi.org/10.1074/jbc.M007694200>
182. Escargueil-Blanc I, Salvayre R, Vacaresse N, Jürgens G, Darblade B, Arnal JF, Parthasarathy S, Nègre-Salvayre A. Mildly oxidized LDL induces activation of platelet-derived growth factor beta-receptor pathway. *Circulation*. 2001; 104:1814–21. <https://doi.org/10.1161/hc4001.097179>
183. Vindis C, Escargueil-Blanc I, Uchida K, Elbaz M, Salvayre R, Negre-Salvayre A. Lipid oxidation products and oxidized low-density lipoproteins impair platelet-derived growth factor receptor activity in smooth muscle cells: implication in atherosclerosis. *Redox Rep*. 2007; 12:96–100. <https://doi.org/10.1179/135100007X162248>
184. Aluise CD, Rose K, Boiani M, Reyzer ML, Manna JD, Tallman K, Porter NA, Marnett LJ. Peptidyl-prolyl cis/trans-isomerase A1 (Pin1) is a target for modification by lipid electrophiles. *Chem Res Toxicol*. 2013; 26:270–79. <https://doi.org/10.1021/tx300449g>
185. Barrera G, Pizzimenti S, Ciamporcero ES, Daga M, Ullio C, Arcaro A, Cetrangolo GP, Ferretti C, Dianzani C, Lepore A, Gentile F. Role of 4-hydroxynonenal-protein adducts in human diseases. *Antioxid Redox Signal*. 2015; 22:1681–702. <https://doi.org/10.1089/ars.2014.6166>
186. Keller JN, Pang Z, Geddes JW, Begley JG, Germeyer A, Waeg G, Mattson MP. Impairment of glucose and glutamate transport and induction of mitochondrial oxidative stress and dysfunction in synaptosomes by amyloid beta-peptide: role of the lipid peroxidation product 4-hydroxynonenal. *J Neurochem*. 1997; 69:273–84. <https://doi.org/10.1046/j.1471-159.1997.69010273.x>
187. Bruce-Keller AJ, Li YJ, Lovell MA, Kraemer PJ, Gary DS, Brown RR, Markesbery WR, Mattson MP. 4-Hydroxynonenal, a product of lipid peroxidation, damages cholinergic neurons and impairs visuospatial memory in rats. *J Neuropathol Exp Neurol*. 1998; 57:257–67. <https://doi.org/10.1097/00005072-199803000-00007>
188. Kruman I, Bruce-Keller AJ, Bredesen D, Waeg G, Mat-

- tson MP. Evidence that 4-hydroxynonenal mediates oxidative stress-induced neuronal apoptosis. *J Neurosci.* 1997; 17:5089–100. <https://doi.org/10.1523/JNEUROSCI.17-13-5089.1997>
189. Nam SM, Chung TH, Kim JW, Jung HY, Yim HS, Kim DW, Yoo DY, Nam H, Choi JH, Hwang IK, Suh JG, Yoon YS. Comparison of N-methyl-D-aspartate receptor subunit 1 and 4-hydroxynonenal in the hippocampus of natural and chemical-induced aging accelerated mice. *Neurochem Res.* 2014; 39:1702–08. <https://doi.org/10.1007/s11064-014-1362-7>
190. Di Domenico F, Tramutola A, Butterfield DA. Role of 4-hydroxy-2-nonenal (HNE) in the pathogenesis of alzheimer disease and other selected age-related neurodegenerative disorders. *Free Radic Biol Med.* 2017; 111:253–61. <https://doi.org/10.1016/j.freeradbiomed.2016.10.490>
191. Cheignon C, Tomas M, Bonnefont-Rousselot D, Faller P, Hureau C, Collin F. Oxidative stress and the amyloid beta peptide in Alzheimer's disease. *Redox Biol.* 2018; 14:450–64. <https://doi.org/10.1016/j.redox.2017.10.014>
192. Hardas SS, Sultana R, Clark AM, Beckett TL, Szweda LI, Murphy MP, Butterfield DA. Oxidative modification of lipoic acid by HNE in Alzheimer disease brain. *Redox Biol.* 2013; 1:80–85. <https://doi.org/10.1016/j.redox.2013.01.002>
193. Sayre LM, Zelasko DA, Harris PL, Perry G, Salomon RG, Smith MA. 4-Hydroxynonenal-derived advanced lipid peroxidation end products are increased in Alzheimer's disease. *J Neurochem.* 1997; 68:2092–97. <https://doi.org/10.1046/j.1471-159.1997.68052092.x>
194. Mark RJ, Pang Z, Geddes JW, Uchida K, Mattson MP. Amyloid beta-peptide impairs glucose transport in hippocampal and cortical neurons: involvement of membrane lipid peroxidation. *J Neurosci.* 1997; 17:1046–54. <https://doi.org/10.1523/JNEUROSCI.17-03-01046.1997>
195. Perluigi M, Sultana R, Cenini G, Di Domenico F, Memo M, Pierce WM, Coccia R, Butterfield DA. Redox proteomics identification of 4-hydroxynonenal-modified brain proteins in Alzheimer's disease: role of lipid peroxidation in Alzheimer's disease pathogenesis. *Proteomics Clin Appl.* 2009; 3:682–93. <https://doi.org/10.1002/prca.200800161>
196. Sultana R, Perluigi M, Allan Butterfield D. Lipid peroxidation triggers neurodegeneration: a redox proteomics view into the Alzheimer disease brain. *Free Radic Biol Med.* 2013; 62:157–69. <https://doi.org/10.1016/j.freeradbiomed.2012.09.027>
197. Mancuso C, Barone E. The heme oxygenase/biliverdin reductase pathway in drug research and development. *Curr Drug Metab.* 2009; 10:579–94. <https://doi.org/10.2174/138920009789375405>
198. Hensley K, Christov A, Kamat S, Zhang XC, Jackson KW, Snow S, Post J. Proteomic identification of binding partners for the brain metabolite lanthionine ketimine (LK) and documentation of LK effects on microglia and motoneuron cell cultures. *J Neurosci.* 2010; 30:2979–88. <https://doi.org/10.1523/JNEUROSCI.5247-09.2010>
199. Pedersen WA, Cashman NR, Mattson MP. The lipid peroxidation product 4-hydroxynonenal impairs glutamate and glucose transport and choline acetyltransferase activity in NSC-19 motor neuron cells. *Exp Neurol.* 1999; 155:1–10. <https://doi.org/10.1006/exnr.1998.6890>
200. Poon HF, Hensley K, Thongboonkerd V, Merchant ML, Lynn BC, Pierce WM, Klein JB, Calabrese V, Butterfield DA. Redox proteomics analysis of oxidatively modified proteins in G93A-SOD1 transgenic mice—a model of familial amyotrophic lateral sclerosis. *Free Radic Biol Med.* 2005; 39:453–62. <https://doi.org/10.1016/j.freeradbiomed.2005.03.030>
201. Kabuta C, Kono K, Wada K, Kabuta T. 4-Hydroxynonenal induces persistent insolubilization of TDP-43 and alters its intracellular localization. *Biochem Biophys Res Commun.* 2015; 463:82–87. <https://doi.org/10.1016/j.bbrc.2015.05.027>
202. Shibata N, Inose Y, Toi S, Hiroi A, Yamamoto T, Kobayashi M. Involvement of 4-hydroxy-2-nonenal accumulation in multiple system atrophy. *Acta Histochem Cytochem.* 2010; 43:69–75. <https://doi.org/10.1267/ahc.10005>
203. Almandoz-Gil L, Welander H, Ihse E, Khoonsari PE, Musunuri S, Lendel C, Sigvardson J, Karlsson M, Ingelsson M, Kultima K, Bergström J. Low molar excess of 4-oxo-2-nonenal and 4-hydroxy-2-nonenal promote oligomerization of alpha-synuclein through different pathways. *Free Radic Biol Med.* 2017. <https://doi.org/10.1016/j.freeradbiomed.2017.07.004>
204. Pecorelli A, Cervellati C, Cortelazzo A, Cervellati F, Sticozzi C, Mirasole C, Guerranti R, Trentini A, Zolla L, Savelli V, Hayek J, Valacchi G. Proteomic analysis of 4-hydroxynonenal and nitrotyrosine modified proteins in RTT fibroblasts. *Int J Biochem Cell Biol.* 2016; 81:236–45. <https://doi.org/10.1016/j.biocel.2016.08.001>
205. Aslebagh R, Pfeffer BA, Fliesler SJ, Darie CC. Mass spectrometry-based proteomics of oxidative stress: identification of 4-hydroxy-2-nonenal (HNE) adducts of amino acids using lysozyme and bovine serum albumin as model proteins. *Electrophoresis.* 2016; 37:2615–23. <https://doi.org/10.1002/elps.201600134>

206. Delosière M, Santé-Lhoutellier V, Chantelauze C, Durand D, Thomas A, Joly C, Pujos-Guillot E, Rémond D, Comte B, Gladine C, Guy A, Durand T, Laurentie M, Dufour C. Quantification of 4-hydroxy-2-nonenal-protein adducts in the in vivo gastric digesta of mini-pigs using a GC-MS/MS method with accuracy profile validation. *Food Funct.* 2016; 7:3497–504. <https://doi.org/10.1039/C6FO00413J>
207. Spickett CM, Wiswedel I, Siems W, Zarkovic K, Zarkovic N. Advances in methods for the determination of biologically relevant lipid peroxidation products. *Free Radic Res.* 2010; 44:1172–202. <https://doi.org/10.3109/10715762.2010.498476>
208. Nag S, Begley DJ. Blood-brain barrier, exchange of metabolites and gases. In: *Pathology and genetics, cerebrovascular diseases*. ISN Neuropath Press, Basel 2005, pp. 22–29.
209. Pan Y, Nicolazzo JA. Impact of aging, Alzheimer's disease and Parkinson's disease on the blood-brain barrier transport of therapeutics. *Adv Drug Deliv Rev.* 2018S0169-409X(18)30061-9.
210. Neuwelt E, Abbott NJ, Abrey L, Banks WA, Blakley B, Davis T, Engelhardt B, Grammas P, Nedergaard M, Nutt J, Pardridge W, Rosenberg GA, Smith Q, Drewes LR. Strategies to advance translational research into brain barriers. *Lancet Neurol.* 2008; 7:84–96. [https://doi.org/10.1016/S1474-4422\(07\)70326-5](https://doi.org/10.1016/S1474-4422(07)70326-5)
211. Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H, Hashimoto N, Furuse M, Tsukita S. Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell Biol.* 2003; 161:653–60. <https://doi.org/10.1083/jcb.200302070>
212. Peters A, Josephson K, Vincent SL. Effects of aging on the neuroglial cells and pericytes within area 17 of the rhesus monkey cerebral cortex. *Anat Rec.* 1991; 229:384–98. <https://doi.org/10.1002/ar.1092290311>
213. Bors L, Tóth K, Tóth EZ, Bajza Á, Csorba A, Szigeti K, Máthé D, Perlaki G, Orsi G, Tóth GK, Erdő F. Age-dependent changes at the blood-brain barrier. A Comparative structural and functional study in young adult and middle aged rats. *Brain Res Bull.* 2018; 139:269–77. <https://doi.org/10.1016/j.brainresbull.2018.03.001>
214. Church RM, Miller MC, Freestone D, Chiu C, Osgood DP, Machan JT, Messier AA, Johanson CE, Silverberg GD. Amyloid-beta accumulation, neurogenesis, behavior, and the age of rats. *Behav Neurosci.* 2014; 128:523–36. <https://doi.org/10.1037/a0036433>
215. Ashraf T, Kis O, Banerjee N, Bendayan R. Drug transporters at brain barriers: expression and regulation by neurological disorders. *Adv Exp Med Biol.* 2012; 763:20–69.
216. van Assema DM, Lubberink M, Boellaard R, Schuit RC, Windhorst AD, Scheltens P, Lammertsma AA, van Berckel BN. P-glycoprotein function at the blood-brain barrier: effects of age and gender. *Mol Imaging Biol.* 2012; 14:771–76. <https://doi.org/10.1007/s11307-012-0556-0>
217. Massaad CA. Neuronal and vascular oxidative stress in Alzheimer's disease. *Curr Neuropharmacol.* 2011; 9:662–73. <https://doi.org/10.2174/157015911798376244>
218. Mertsch K, Blasig I, Grune T. 4-Hydroxynonenal impairs the permeability of an in vitro rat blood-brain barrier. *Neurosci Lett.* 2001; 314:135–38. [https://doi.org/10.1016/S0304-3940\(01\)02299-6](https://doi.org/10.1016/S0304-3940(01)02299-6)
219. Wang J, Sun L, Si YF, Li BM. Overexpression of actin-depolymerizing factor blocks oxidized low-density lipoprotein-induced mouse brain microvascular endothelial cell barrier dysfunction. *Mol Cell Biochem.* 2012; 371:1–8. <https://doi.org/10.1007/s11010-012-1415-7>
220. Schommer J, Marwarha G, Schommer T, Flick T, Lund J, Ghribi O. 27-Hydroxycholesterol increases α -synuclein protein levels through proteasomal inhibition in human dopaminergic neurons. *BMC Neurosci.* 2018; 19:17. <https://doi.org/10.1186/s12868-018-0420-5>
221. Powers ET, Morimoto RI, Dillin A, Kelly JW, Balch WE. Biological and chemical approaches to diseases of proteostasis deficiency. *Annu Rev Biochem.* 2009; 78:959–91. <https://doi.org/10.1146/annurev.biochem.052308.114844>
222. Hartl FU. Chaperone-assisted protein folding: the path to discovery from a personal perspective. *Nat Med.* 2011; 17:1206–10. <https://doi.org/10.1038/nm.2467>
223. Koga H, Kaushik S, Cuervo AM. Protein homeostasis and aging: the importance of exquisite quality control. *Ageing Res Rev.* 2011; 10:205–15. <https://doi.org/10.1016/j.arr.2010.02.001>
224. van Ham TJ, Holmberg MA, van der Goot AT, Teuling E, Garcia-Arencibia M, Kim HE, Du D, Thijssen KL, Wiersma M, Burggraaff R, van Bergeijk P, van Rheenen J, Jerre van Veluw G, et al. Identification of MOAG-4/SERF as a regulator of age-related proteotoxicity. *Cell.* 2010; 142:601–12. <https://doi.org/10.1016/j.cell.2010.07.020>
225. Chondrogianni N, Gonos ES. Proteasome function determines cellular homeostasis and the rate of aging. *Adv Exp Med Biol.* 2010; 694:38–46. https://doi.org/10.1007/978-1-4419-7002-2_4

226. Chondrogianni N, Voutetakis K, Kapetanou M, Delitsikou V, Papaevgeniou N, Sakellari M, Lefaki M, Filippopoulou K, Gonos ES. Proteasome activation: an innovative promising approach for delaying aging and retarding age-related diseases. *Ageing Res Rev.* 2015; 23:37–55. <https://doi.org/10.1016/j.arr.2014.12.003>
227. Goldberg AL. Functions of the proteasome: from protein degradation and immune surveillance to cancer therapy. *Biochem Soc Trans.* 2007; 35:12–17. <https://doi.org/10.1042/BST0350012>
228. Chondrogianni N, Georgila K, Kourtis N, Tavernarakis N, Gonos ES. 20S proteasome activation promotes life span extension and resistance to proteotoxicity in *Caenorhabditis elegans*. *FASEB J.* 2015; 29:611–22. <https://doi.org/10.1096/fj.14-252189>
229. da Fonseca PC, Morris EP. Structure of the human 26S proteasome: subunit radial displacements open the gate into the proteolytic core. *J Biol Chem.* 2008; 283:23305–14. <https://doi.org/10.1074/jbc.M802716200>
230. Tanaka K. The proteasome: from basic mechanisms to emerging roles. *Keio J Med.* 2013; 62:1–12. <https://doi.org/10.2302/kjm.2012-0006-RE>
231. Nickell S, Beck F, Scheres SH, Korinek A, Förster F, Lasker K, Mihalache O, Sun N, Nagy I, Sali A, Plitzko JM, Carazo JM, Mann M, Baumeister W. Insights into the molecular architecture of the 26S proteasome. *Proc Natl Acad Sci USA.* 2009; 106:11943–47. <https://doi.org/10.1073/pnas.0905081106>
232. Koga H, Martinez-Vicente M, Arias E, Kaushik S, Sulzer D, Cuervo AM. Constitutive upregulation of chaperone-mediated autophagy in Huntington's disease. *J Neurosci.* 2011; 31:18492–505. <https://doi.org/10.1523/JNEUROSCI.3219-11.2011>
233. Bucciantini M, Giannoni E, Chiti F, Baroni F, Formigli L, Zurdo J, Taddei N, Ramponi G, Dobson CM, Stefani M. Inherent toxicity of aggregates implies a common mechanism for protein misfolding diseases. *Nature.* 2002; 416:507–11. <https://doi.org/10.1038/416507a>
234. Jana NR. Protein homeostasis and aging: role of ubiquitin protein ligases. *Neurochem Int.* 2012; 60:443–47. <https://doi.org/10.1016/j.neuint.2012.02.009>
235. Chondrogianni N, Sakellari M, Lefaki M, Papaevgeniou N, Gonos ES. Proteasome activation delays aging in vitro and in vivo. *Free Radic Biol Med.* 2014; 71:303–20. <https://doi.org/10.1016/j.freeradbiomed.2014.03.031>
236. Chondrogianni N, Tzavelas C, Pemberton AJ, Nezis IP, Rivett AJ, Gonos ES. Overexpression of proteasome beta5 assembled subunit increases the amount of proteasome and confers ameliorated response to oxidative stress and higher survival rates. *J Biol Chem.* 2005; 280:11840–50. <https://doi.org/10.1074/jbc.M413007200>
237. Chondrogianni N, Trougakos IP, Kletsas D, Chen QM, Gonos ES. Partial proteasome inhibition in human fibroblasts triggers accelerated M1 senescence or M2 crisis depending on p53 and Rb status. *Aging Cell.* 2008; 7:717–32. <https://doi.org/10.1111/j.1474-9726.2008.00425.x>
238. Grune T, Jung T, Merker K, Davies KJ. Decreased proteolysis caused by protein aggregates, inclusion bodies, plaques, lipofuscin, ceroid, and 'aggresomes' during oxidative stress, aging, and disease. *Int J Biochem Cell Biol.* 2004; 36:2519–30. <https://doi.org/10.1016/j.biocel.2004.04.020>
239. Andersson V, Hanzén S, Liu B, Molin M, Nyström T. Enhancing protein disaggregation restores proteasome activity in aged cells. *Aging (Albany NY).* 2013; 5:802–12. <https://doi.org/10.18632/aging.100613>
240. Tonoki A, Kuranaga E, Tomioka T, Hamazaki J, Murata S, Tanaka K, Miura M. Genetic evidence linking age-dependent attenuation of the 26S proteasome with the aging process. *Mol Cell Biol.* 2009; 29:1095–106. <https://doi.org/10.1128/MCB.01227-08>
241. Vernace VA, Arnaud L, Schmidt-Glenewinkel T, Figueiredo-Pereira ME. Aging perturbs 26S proteasome assembly in *Drosophila melanogaster*. *FASEB J.* 2007; 21:2672–82. <https://doi.org/10.1096/fj.06-6751com>
242. Rodriguez KA, Edrey YH, Osmulski P, Gaczynska M, Buffenstein R. Altered composition of liver proteasome assemblies contributes to enhanced proteasome activity in the exceptionally long-lived naked mole-rat. *PLoS One.* 2012; 7:e35890. <https://doi.org/10.1371/journal.pone.0035890>
243. Chondrogianni N, Petropoulos I, Franceschi C, Friguet B, Gonos ES. Fibroblast cultures from healthy centenarians have an active proteasome. *Exp Gerontol.* 2000; 35:721–28. [https://doi.org/10.1016/S0531-5565\(00\)00137-6](https://doi.org/10.1016/S0531-5565(00)00137-6)
244. Vilchez D, Boyer L, Morantte I, Lutz M, Merkwirth C, Joyce D, Spencer B, Page L, Masliah E, Berggren WT, Gage FH, Dillin A. Increased proteasome activity in human embryonic stem cells is regulated by PSMD11. *Nature.* 2012; 489:304–08. <https://doi.org/10.1038/nature11468>
245. Kapetanou M, Chondrogianni N, Petrakis S, Koliakos G, Gonos ES. Proteasome activation enhances stemness and lifespan of human mesenchymal stem cells. *Free Radic Biol Med.* 2017; 103:226–35. <https://doi.org/10.1016/j.freeradbiomed.2016.12.035>

246. Zhang C, Cuervo AM. Restoration of chaperone-mediated autophagy in aging liver improves cellular maintenance and hepatic function. *Nat Med*. 2008; 14:959–65. <https://doi.org/10.1038/nm.1851>
247. Liu G, Rogers J, Murphy CT, Rongo C. EGF signalling activates the ubiquitin proteasome system to modulate *C. elegans* lifespan. *EMBO J*. 2011; 30:2990–3003. <https://doi.org/10.1038/emboj.2011.195>
248. Kruegel U, Robison B, Dange T, Kahlert G, Delaney JR, Kotireddy S, Tsuchiya M, Tsuchiyama S, Murakami CJ, Schleit J, Sutphin G, Carr D, Tar K, et al. Elevated proteasome capacity extends replicative lifespan in *Saccharomyces cerevisiae*. *PLoS Genet*. 2011; 7:e1002253. <https://doi.org/10.1371/journal.pgen.1002253>
249. Vilchez D, Morante I, Liu Z, Douglas PM, Merkwirth C, Rodrigues AP, Manning G, Dillin A. RPN-6 determines *C. elegans* longevity under proteotoxic stress conditions. *Nature*. 2012; 489:263–68. <https://doi.org/10.1038/nature11315>
250. Hwang JS, Hwang JS, Chang I, Kim S. Age-associated decrease in proteasome content and activities in human dermal fibroblasts: restoration of normal level of proteasome subunits reduces aging markers in fibroblasts from elderly persons. *J Gerontol A Biol Sci Med Sci*. 2007; 62:490–99. <https://doi.org/10.1093/gerona/62.5.490>
251. Liu Y, Liu X, Zhang T, Luna C, Liton PB, Gonzalez P. Cytoprotective effects of proteasome beta5 subunit overexpression in lens epithelial cells. *Mol Vis*. 2007; 13:31–38.
252. Malhotra D, Thimmulappa R, Vij N, Navas-Acien A, Sussan T, Merali S, Zhang L, Kelsen SG, Myers A, Wise R, Tudor R, Biswal S. Heightened endoplasmic reticulum stress in the lungs of patients with chronic obstructive pulmonary disease: the role of Nrf2-regulated proteasomal activity. *Am J Respir Crit Care Med*. 2009; 180:1196–207. <https://doi.org/10.1164/rccm.200903-0324OC>
253. Chondrogianni N, Gonos ES. Overexpression of hUMP1/POMP proteasome accessory protein enhances proteasome-mediated antioxidant defence. *Exp Gerontol*. 2007; 42:899–903. <https://doi.org/10.1016/j.exger.2007.01.012>
254. Li J, Horak KM, Su H, Sanbe A, Robbins J, Wang X. Enhancement of proteasomal function protects against cardiac proteinopathy and ischemia/reperfusion injury in mice. *J Clin Invest*. 2011; 121:3689–700. <https://doi.org/10.1172/JCI45709>
255. Campos MG, Webby RF, Markham KR, Mitchell KA, Da Cunha AP. Age-induced diminution of free radical scavenging capacity in bee pollens and the contribution of constituent flavonoids. *J Agric Food Chem*. 2003; 51:742–45. <https://doi.org/10.1021/jf0206466>
256. Graikou K, Kapeta S, Aligiannis N, Sotiroudis G, Chondrogianni N, Gonos E, Chinou I. Chemical analysis of Greek pollen - Antioxidant, antimicrobial and proteasome activation properties. *Chem Cent J*. 2011; 5:33. <https://doi.org/10.1186/1752-153X-5-33>
257. Katsiki M, Chondrogianni N, Chinou I, Rivett AJ, Gonos ES. The olive constituent oleuropein exhibits proteasome stimulatory properties in vitro and confers life span extension of human embryonic fibroblasts. *Rejuvenation Res*. 2007; 10:157–72. <https://doi.org/10.1089/rej.2006.0513>
258. Ali RE, Rattan SI. Curcumin's biphasic hormetic response on proteasome activity and heat-shock protein synthesis in human keratinocytes. *Ann N Y Acad Sci*. 2006; 1067:394–99. <https://doi.org/10.1196/annals.1354.056>
259. Dal Vechio FH, Cerqueira F, Augusto O, Lopes R, Demasi M. Peptides that activate the 20S proteasome by gate opening increased oxidized protein removal and reduced protein aggregation. *Free Radic Biol Med*. 2014; 67:304–13. <https://doi.org/10.1016/j.freeradbiomed.2013.11.017>
260. Kwak MK, Cho JM, Huang B, Shin S, Kensler TW. Role of increased expression of the proteasome in the protective effects of sulforaphane against hydrogen peroxide-mediated cytotoxicity in murine neuroblastoma cells. *Free Radic Biol Med*. 2007; 43:809–17. <https://doi.org/10.1016/j.freeradbiomed.2007.05.029>
261. Kapeta S, Chondrogianni N, Gonos ES. Nuclear erythroid factor 2-mediated proteasome activation delays senescence in human fibroblasts. *J Biol Chem*. 2010; 285:8171–84. <https://doi.org/10.1074/jbc.M109.031575>
262. Papaevgeniou N, Sakellari M, Jha S, Tavernarakis N, Holmberg CI, Gonos ES, Chondrogianni N. 18 α -Glycyrrhetic Acid Proteasome Activator Decelerates Aging and Alzheimer's Disease Progression in *Caenorhabditis elegans* and Neuronal Cultures. *Antioxid Redox Signal*. 2016; 25:855–69. <https://doi.org/10.1089/ars.2015.6494>
263. Chondrogianni N, Kapeta S, Chinou I, Vassilatou K, Papassideri I, Gonos ES. Anti-ageing and rejuvenating effects of quercetin. *Exp Gerontol*. 2010; 45:763–71. <https://doi.org/10.1016/j.exger.2010.07.001>
264. Jang J, Wang Y, Kim HS, Lalli MA, Kosik KS. Nrf2, a regulator of the proteasome, controls self-renewal and pluripotency in human embryonic stem cells. *Stem Cells*. 2014; 32:2616–25. <https://doi.org/10.1002/stem.1764>