

Brown adipose tissue enhances exercise performance and healthful longevity

Dorothy E. Vatner¹, Jie Zhang¹, Stephen F. Vatner¹

¹Department of Cell Biology and Molecular Medicine, Rutgers New Jersey Medical School, Newark, NJ 07103, USA

Correspondence to: Stephen F. Vatner; email: vatnersf@njms.rutgers.edu

Keywords: brown adipose tissue, white adipose tissue, healthful longevity, exercise, regulator of G protein signaling 14

Received: November 13, 2024

Accepted: December 2, 2024

Published: December 18, 2024

Copyright: © 2024 Vatner et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/) (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Brown adipose tissue (BAT), a major subtypes of adipose tissues, is known for thermogenesis and promoting healthful longevity. Our hypothesis is that BAT protects against impaired healthful longevity, i.e., obesity, diabetes, cardiovascular disorders, cancer, Alzheimer's disease, and reduced exercise tolerance. While most prior studies have shown that exercise regulates BAT activation and improves BAT density, relatively few have shown that BAT increases exercise performance. In contrast, our recent studies with the regulator of G protein signaling 14 (RGS14) knockout (KO) model of extended longevity showed that it enhances exercise performance, mediated by its more potent BAT, compared with BAT from wild type mice. For example, when the BAT from RGS14 KO mice is transplanted to WT mice, their exercise capacity is enhanced at 3 days after BAT transplantation, whereas BAT transplantation from WT to WT mice increased exercise performance, but only at 8 weeks after transplantation. The goal of this research perspective is to review the role of BAT in mediating healthful longevity, specifically exercise capacity. In view of the ability of BAT to mediate healthful longevity and enhance exercise performance, it is likely that a pharmaceutical analog of BAT will become a novel therapeutic modality.

White adipose tissue (WAT) vs. brown adipose tissue (BAT)

There are two major subtypes of adipose tissue, i.e., WAT and BAT, each with distinct functions, cellular structures, and origins. WAT originates from mesenchymal stem cells during development and consists of large, unilocular adipocytes. These cells contain a single lipid droplet and are found in areas like the abdomen, thighs, and around internal organs [1]. WAT functions primarily as an energy reservoir, storing excess calories in the form of triglycerides, but also plays a role in impairing healthful longevity by increasing obesity and its related metabolic disorders, such as insulin resistance, type 2 diabetes, and cardiovascular diseases [2, 3].

Brown adipose tissue (BAT)

In marked contrast to WAT, BAT is becoming increasingly attributed to healthful aging. Brown

adipocytes are smaller in size than white adipocytes with lipid droplets surrounding the nucleus. Brown adipocytes have mitochondria dispersed between the droplets, which give these cells their brown appearance. BAT is mainly located in the interscapular space of mice, and in humans it is found in the interscapular, supraclavicular, suprarenal, and para-aortic spaces. BAT originates from a lineage closely related to skeletal muscle cells, also derived from mesenchymal stem cells. BAT consists of smaller adipocytes with mitochondria, which produce heat through uncoupling protein 1 (UCP1) [4], resulting in the release of heat [1, 5–8]. This provides the basis for the major role for BAT, thermogenesis, as a mechanism to maintain body temperature by helping maintain body temperature in cold environments. This function is especially critical in newborns, who have a higher proportion of BAT, but it persists in adults in smaller amounts, especially around the neck and shoulders [9]. Changes in BAT with aging reduce its effects on thermogenesis [10]. BAT is

activated by norepinephrine, a hormone released during cold exposure and sympathetic nervous system activation. Norepinephrine binds to β -adrenergic receptors on brown adipocytes, stimulating the production of UCP1, which uncouples oxidative phosphorylation from ATP synthesis, allowing energy to be dissipated as heat [5]. Additionally, thyroid hormones are important for BAT activation and thermogenesis, enhancing the tissue's ability to burn fat [11].

In addition, UCP1-independent thermogenic pathways have been found in beige adipocytes and muscles [12]. BAT also functions as a metabolic sink by oxidizing glucose and lipids, which produces heat. Since BAT burns calories to produce heat, its activation promotes weight loss, and enhances insulin sensitivity [13], which have therapeutic potential for obesity and diabetes [14]. BAT has also been shown to mediate healthful longevity [8, 15].

BAT mediating healthful longevity

Studies have indicated that aging reduces BAT activity leading to thermal dysregulation and energy imbalance [16–18]. However, the effects of age on BAT mass have been inconsistent [19–21]. Some studies have reported that aging increases the amount of BAT [19, 20], whereas one study reported no change in BAT mass in rodents [21]. In addition, beige adipocyte formation declines with aging which may be caused by changes in the adipose tissue microenvironment [16, 22].

More recently, interest has extended to BAT's role in mediating healthful aging, primarily from data in genetically altered mouse models. It has been reported that the enhanced BAT activity/function mediate healthful longevity in several longevity mouse models, e.g., protection against obesity [23, 24], diabetes [23, 24], cardiovascular disorders [25–27], cancer [28–32], Alzheimer's Disease [33], stroke [34, 35], exercise intolerance [36], and reduced blood flow [36–38].

In addition, there are models with enhanced BAT function or extra BAT amount by BAT transplantation that exhibit aspects of healthful longevity. WT mice receiving BAT from another WT BAT mouse exhibit improved exercise capacity, protection against obesity and diabetes and cancer [36, 39–41]. It has also been suggested that low levels of BAT in humans are associated with obesity and glucose intolerance, whereas those with higher BAT levels maintain lower body weight and more healthful aging [42].

One example of a genetic model demonstrating that BAT enhances exercise performance is that of RGS14

KO mice, a healthful lifespan model, mediated by increased BAT [8]. Survival is significantly enhanced in the RGS14 KO mouse, with females living longer than males, similar to human data (Figure 1A, 1B). In addition, the old RGS14 KO mice do not show the phenotype of aging WT mice, i.e., body atrophy, loss of hair and greying of hair color (Figure 1C). These features of healthy aging can be recapitulated in WT mice with transplants of BAT from RGS14 KO mice at a young age (Figure 1C). The RGS14 KO mice have increased density of BAT with smaller BAT cell size (Figure 2).

BAT and exercise

Exercise regulating BAT

Enhanced exercise capacity is not only a feature of healthful aging, but also is a therapy for aging patients and patients with cardiovascular disease. Exercise is a healthy way to reduce body weight by activating the sympathetic nervous system, accelerating the decomposition of fat, and promoting the utilization and consumption of energy in skeletal muscle [43–45]. During aging, it is known that progressive loss of exercise capacity relates to loss of skeletal muscle mass and tissue function [46]. Decreased muscle mitochondrial function contributes to the loss of skeletal muscle function during aging [47–51]. Regular exercise or exercise training protects against decreased muscle function during aging [52, 53], frailty status [54, 55], and neurodegeneration [56, 57].

Numerous studies have suggested that exercise may play a role in regulating BAT activation. Exercise boosts the expression of UCP1 and genes associated with mitochondria biogenesis, thereby improving BAT's heat production capacity [5]. For instance, swim training in rodents over six to eight weeks increased UCP1 protein levels in BAT [58, 59]. Similarly, treadmill exercise in rodents for 6–8 weeks increased BAT activity and cytochrome oxidase activity, oxygen consumption rates and BAT-specific gene markers, e.g., UCP1, FGF21, and PGC1 α [60, 61]. However, conflicting findings also exist, with some studies suggesting that exercise may reduce the thermogenic effect of BAT. In rats, six to eight weeks of moderate-intensity treadmill exercise led to decreased UCP1 expression in BAT and a reduction in total BAT mass [62, 63]. Human studies also showed inconsistent results regarding the role of exercise on BAT modulation, with some indicating that high-intensity physical activities can increase BAT density [64], while others reporting that exercise decreases glucose uptake in BAT [65–67]. While most of these studies have shown that exercise increases BAT, relatively few have




Figure 1. RGS14 KO model of longevity. (A) Kaplan-Meier survival curves for RGS14 KO and WT mice showed significantly augmented survival in RGS14 KO mice. (B) Maximum lifespan was significantly greater in RGS14 KO mice than in WT mice for both males and females. In addition, median and maximum lifespan were greater in female RGS14 KO mice than in male RGS14 KO mice. (C) Furthermore, 24 month old RGS14 KO mice did not show the aging phenotype normally present in WT mice of similar age, including body atrophy, loss of hair and greying of fur color. In support of the key role of BAT in aging, old WT RGS14 KO BAT recipient mice, which had BAT transplanted at 3–4 months of age, had the appearance of youthful aging similar to the old RGS14 KO mice. A representative example of each is shown in panel C. For median lifespan analysis, a Mood's Median Test, was used to determine differences in median lifespan. A Student's *t*-test was used to test differences in maximum lifespan. **p* < 0.05 vs. WT, #*p* < 0.05 vs. male. Reprinted from Vatner DE, et al Aging Cell. 2018; 17(4).

RGS14 KO Brown Adipose Tissue




Figure 2. Increased BAT cell numbers in RGS14 KO mice. RGS14 KO mice exhibited smaller brown adipocytes (A, B), and increased number of brown adipocytes (A, C) than WT control mice. **p* < 0.05. Reprinted from Vatner DE, et al Aging Cell. 2023; 22(4).

shown that BAT increases exercise performance [36, 41, 68].

BAT regulating exercise

There are fewer studies examining the role of BAT regulating exercise than exercise regulating BAT. One example of a genetic model demonstrating that BAT can enhance exercise performance is that of RGS14 KO mice, a healthful lifespan model, mediated by increased BAT [8]. Whereas the RGS14 KO mouse exhibits increased exercise capacity, similarly, RGS14 KO BAT transplanted to WT mice demonstrate an increased maximal running distance and work to exhaustion, which corresponds to the enhanced exercise capacity of RGS14 KO mice (Figure 3A–3D). The enhanced exercise capacity observed in WT mice with RGS14

KO BAT transplants was observed at three days after BAT transplantation, whereas BAT transplantation from WT to WT mice also resulted in increased exercise performance, not at 3 days, but only at 8 weeks after transplantation [36] (Figure 3E–3H). Another study found that BAT transplantation prevents the impaired glucose tolerance, the increase in left ventricle mass, and exercise intolerance following myocardial infarction [41]. In addition, BAT transplantation significantly reversed the reduction in physical activity in high fat diet-fed mice [68].

There are multiple mechanisms mediating the enhanced exercise capacity in RGS14 KO mice. Most importantly, RGS14 KO mice demonstrate increased hind limb perfusion (Figure 4A), also found in RGS14 KO BAT recipients, but lost in RGS14 KO BAT donors




Figure 3. Increased exercise capacity in RGS14 KO mice. RGS14 KO mice ran longer distances (A) with increased work to exhaustion (B) compared to WT littermates. BAT transplantation from RGS14 KO mice to WT mice led to a reversal of phenotype, such that RGS14 KO BAT recipients exhibited improved running distance (C) and greater work to exhaustion (D) compared to RGS14 KO BAT donors, at 3 days after RGS14 KO BAT transplantation. In contrast, there was no improvement in running distance and work to exhaustion at 3 days after transplantation of BAT from C57BL6/J WT mice to other C57BL6/J WT mice (E, F). It required 8 weeks to achieve enhanced running distance and work to exhaustion in C57BL6/J WT mice with BAT transplantation from other C57BL6/J WT mice (G, H). * $p < 0.05$. Reprinted from Vatner DE, et al Aging Cell. 2023; 22(4).

Enhanced Exercise in RGS14 KO Mice Is Mediated By Brown Adipose Tissue and Angiogenesis/ Arteriogenesis Increasing Hind Limb Blood Flow




Figure 4. Enhanced exercise by RGS14 KO mice is mediated by BAT and angiogenesis/arteriogenesis increasing hindlimb blood flow. Non-linear contrast imaging was used to measure hindlimb blood flow. The average data are presented as % of WT perfusion, which is represented as 100% (A). Hindlimb blood flow was higher in RGS14 KO mice compared to WT mice, and higher in WT mice that received RGS14 KO BAT, at 3 days after transplantation (A, B), while RGS14 KO BAT donors lost their enhanced hindlimb perfusion, with results similar to WT mice (A). WT BAT recipients showed greater hindlimb perfusion at 8 weeks after transplantation of BAT from C57BL6/J WT mice to other C57BL6/J WT mice (A). Angiogenesis (reflected by capillary density) and arteriogenesis (reflected by arteriole density) were both increased in skeletal muscle of RGS14 KO mice (B, C) and RGS14 KO BAT recipients (D, E), which correlated with increased VEGF in skeletal muscle (F) and BAT (G). Increased angiogenesis (D) and arteriogenesis (E) were not observed in RGS14 KO BAT donors. Results are expressed as Mean $\pm$ SEM, * p < 0.05. Reprinted from Vatner DE, et al Aging Cell. 2023; 22(4).




Figure 5. Mechanisms mediating enhanced exercise capacity in RGS14 KO and its uniquely powerful BAT. Multiple mechanisms mediated the enhanced exercise capacity in RGS14 KO mice. The most important mechanism is BAT, which mediates SIRT3, MnSOD, MEK/ERK and VEGF pathways. These mechanisms regulate exercise capacity by improved mitochondrial function, protection against oxidative stress and improved blood flow/angiogenesis. Reprinted from Vatner DE, et al Aging Cell. 2023; 22(4).

(Figure 4A) and increased capillary and arteriolar density (Figure 4B, 4C). The increase capillary and arteriolar density was also observed in RGS14 KO BAT recipients and lost in RGS14 KO BAT donors (Figure 4D, 4E). A key mechanism mediating the increased perfusion and vascular density was VEGF (Figure 4F), observed in RGS14 KO skeletal muscle and BAT (Figure 4G). At the cellular level, the mechanisms mediating the RGS14 KO and RGS14 KO BAT enhanced exercise capacity include increases in SIRT3, MEK/ERK, MnSOD, and mitochondrial function, which decrease oxidative stress. All these mechanisms are illustrated in Figure 5.

In view of the data demonstrating that the RGS14 KO mouse exhibits many features of healthful aging, one of which is enhanced exercise capacity, mediated by BAT, and that RGS14 KO BAT is more powerful than WT BAT, it becomes increasingly important to develop a pharmacological analog of RGS14 KO BAT that can be translated to the clinics to promote enhanced exercise capacity and healthful aging in patients.

AUTHOR CONTRIBUTIONS

Design of Review: S.F.V.; D.E.V. Writing of the Manuscript: S.F.V.; D.E.V.; J.Z.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest related to this study.

FUNDING

This study was supported by National Institutes of Health grant R21AG075656 (to S.F.V.).

REFERENCES

- Rosen ED, Spiegelman BM. What we talk about when we talk about fat. *Cell*. 2014; 156:20–44. <https://doi.org/10.1016/j.cell.2013.12.012> PMID:24439368
- Galic S, Oakhill JS, Steinberg GR. Adipose tissue as an endocrine organ. *Mol Cell Endocrinol*. 2010; 316:129–39. <https://doi.org/10.1016/j.mce.2009.08.018> PMID:19723556
- Ouchi N, Parker JL, Lugus JJ, Walsh K. Adipokines in inflammation and metabolic disease. *Nat Rev Immunol*. 2011; 11:85–97. <https://doi.org/10.1038/nri2921> PMID:21252989
- Seale P, Bjork B, Yang W, Kajimura S, Chin S, Kuang S, Scimè A, Devarakonda S, Conroe HM, Erdjument-Bromage H, Tempst P, Rudnicki MA, Beier DR, Spiegelman BM. PRDM16 controls a brown fat/skeletal muscle switch. *Nature*. 2008; 454:961–7. <https://doi.org/10.1038/nature07182> PMID:18719582
- Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiol Rev*. 2004; 84:277–359. <https://doi.org/10.1152/physrev.00015.2003> PMID:14715917
- Himms-Hagen J. Brown adipose tissue thermogenesis and obesity. *Prog Lipid Res*. 1989; 28:67–115. [https://doi.org/10.1016/0163-7827\(89\)90009-x](https://doi.org/10.1016/0163-7827(89)90009-x) PMID:2692032
- Li X, Frazier JA, Spahiu E, McPherson M, Miller RA. Muscle-dependent regulation of adipose tissue function in long-lived growth hormone-mutant mice. *Aging (Albany NY)*. 2020; 12:8766–89. <https://doi.org/10.18632/aging.103380> PMID:32464603
- Vatner DE, Zhang J, Oydanich M, Guers J, Katsyuba E, Yan L, Sinclair D, Auwerx J, Vatner SF. Enhanced longevity and metabolism by brown adipose tissue with disruption of the regulator of G protein signaling 14. *Aging Cell*. 2018; 17:e12751. <https://doi.org/10.1111/acer.12751> PMID:29654651
- Cypess AM, Lehman S, Williams G, Tal I, Rodman D, Goldfine AB, Kuo FC, Palmer EL, Tseng YH, Doria A, Kolodny GM, Kahn CR. Identification and importance of brown adipose tissue in adult humans. *N Engl J Med*. 2009; 360:1509–17. <https://doi.org/10.1056/NEJMoa0810780> PMID:19357406
- Gohlke S, Zagoriy V, Cuadros Inostroza A, Méret M, Mancini C, Japtok L, Schumacher F, Kuhlow D, Graja A, Stephanowitz H, Jähnert M, Krause E, Wernitz A, et al. Identification of functional lipid metabolism biomarkers of brown adipose tissue aging. *Mol Metab*. 2019; 24:1–17. <https://doi.org/10.1016/j.molmet.2019.03.011> PMID:31003944
- Silva JE. Thermogenic mechanisms and their hormonal regulation. *Physiol Rev*. 2006; 86:435–64. <https://doi.org/10.1152/physrev.00009.2005> PMID:16601266
- Ikeda K, Yamada T. UCP1 Dependent and Independent Thermogenesis in Brown and Beige Adipocytes. *Front Endocrinol (Lausanne)*. 2020; 11:498.

<https://doi.org/10.3389/fendo.2020.00498>

PMID:[32849287](#)

13. Chondronikola M, Volpi E, Børnsheim E, Porter C, Annamalai P, Enerbäck S, Lidell ME, Saraf MK, Labbe SM, Hurren NM, Yfanti C, Chao T, Andersen CR, et al. Brown adipose tissue improves whole-body glucose homeostasis and insulin sensitivity in humans. *Diabetes*. 2014; 63:4089–99.
<https://doi.org/10.2337/db14-0746>
PMID:[25056438](#)
14. Betz MJ, Enerbäck S. Human Brown Adipose Tissue: What We Have Learned So Far. *Diabetes*. 2015; 64:2352–60.
<https://doi.org/10.2337/db15-0146>
PMID:[26050667](#)
15. Zhang J, Kibret BG, Vatner DE, Vatner SF. The role of brown adipose tissue in mediating healthful longevity. *J Cardiovasc Aging*. 2024; 4:17.
<https://doi.org/10.20517/jca.2024.01>
PMID:[39119146](#)
16. Zoico E, Rubele S, De Caro A, Nori N, Mazzali G, Fantin F, Rossi A, Zamboni M. Brown and Beige Adipose Tissue and Aging. *Front Endocrinol (Lausanne)*. 2019; 10:368.
<https://doi.org/10.3389/fendo.2019.00368>
PMID:[31281288](#)
17. Berry DC, Jiang Y, Arpke RW, Close EL, Uchida A, Reading D, Berglund ED, Kyba M, Graff JM. Cellular Aging Contributes to Failure of Cold-Induced Beige Adipocyte Formation in Old Mice and Humans. *Cell Metab*. 2017; 25:166–81.
<https://doi.org/10.1016/j.cmet.2016.10.023>
PMID:[27889388](#)
18. Lee P, Swarbrick MM, Ho KK. Brown adipose tissue in adult humans: a metabolic renaissance. *Endocr Rev*. 2013; 34:413–38.
<https://doi.org/10.1210/er.2012-1081>
PMID:[23550082](#)
19. Félix-Soriano E, Sáinz N, Gil-Iturbe E, Collantes M, Fernández-Galilea M, Castilla-Madriral R, Ly L, Dalli J, Moreno-Aliaga MJ. Changes in brown adipose tissue lipid mediator signatures with aging, obesity, and DHA supplementation in female mice. *FASEB J*. 2021; 35:e21592.
<https://doi.org/10.1096/fj.202002531R>
PMID:[33960028](#)
20. Gonçalves LF, Machado TQ, Castro-Pinheiro C, de Souza NG, Oliveira KJ, Fernandes-Santos C. Ageing is associated with brown adipose tissue remodelling and loss of white fat browning in female C57BL/6 mice. *Int J Exp Pathol*. 2017; 98:100–8.
<https://doi.org/10.1111/iep.12228>
PMID:[28543963](#)
21. Sellayah D, Sikder D. Orexin restores aging-related brown adipose tissue dysfunction in male mice. *Endocrinology*. 2014; 155:485–501.
<https://doi.org/10.1210/en.2013-1629>
PMID:[24248466](#)
22. Ou MY, Zhang H, Tan PC, Zhou SB, Li QF. Adipose tissue aging: mechanisms and therapeutic implications. *Cell Death Dis*. 2022; 13:300.
<https://doi.org/10.1038/s41419-022-04752-6>
PMID:[35379822](#)
23. Saito M, Okamatsu-Ogura Y. Thermogenic Brown Fat in Humans: Implications in Energy Homeostasis, Obesity and Metabolic Disorders. *World J Mens Health*. 2023; 41:489–507.
<https://doi.org/10.5534/wjmh.220224>
PMID:[36792089](#)
24. Yuko OO, Saito M. Brown Fat as a Regulator of Systemic Metabolism beyond Thermogenesis. *Diabetes Metab J*. 2021; 45:840–52.
<https://doi.org/10.4093/dmj.2020.0291>
PMID:[34176254](#)
25. Becher T, Palanisamy S, Kramer DJ, Eljalby M, Marx SJ, Wibmer AG, Butler SD, Jiang CS, Vaughan R, Schöder H, Mark A, Cohen P. Brown adipose tissue is associated with cardiometabolic health. *Nat Med*. 2021; 27:58–65.
<https://doi.org/10.1038/s41591-020-1126-7>
PMID:[33398160](#)
26. Pereira RO, McFarlane SI. The Role of Brown Adipose Tissue in Cardiovascular Disease Protection: Current Evidence and Future Directions. *Int J Clin Res Trials*. 2019; 4:136.
<https://doi.org/10.15344/2456-8007/2019/136>
PMID:[31650091](#)
27. Chen HJ, Meng T, Gao PJ, Ruan CC. The Role of Brown Adipose Tissue Dysfunction in the Development of Cardiovascular Disease. *Front Endocrinol (Lausanne)*. 2021; 12:652246.
<https://doi.org/10.3389/fendo.2021.652246>
PMID:[34113316](#)
28. Tsoi M, Moore M, Burg D, Painter A, Taylor R, Lockie SH, Turner N, Warren A, Cooney G, Oldfield B, Clarke S, Robertson G. Activation of thermogenesis in brown adipose tissue and dysregulated lipid metabolism associated with cancer cachexia in mice. *Cancer Res*. 2012; 72:4372–82.
<https://doi.org/10.1158/0008-5472.CAN-11-3536>
PMID:[22719069](#)
29. Cao Q, Hersl J, La H, Smith M, Jenkins J, Goloubeva O, Dilszian V, Tkaczuk K, Chen W, Jones L. A pilot study of FDG PET/CT detects a link between brown adipose

- tissue and breast cancer. *BMC Cancer*. 2014; 14:126.
<https://doi.org/10.1186/1471-2407-14-126>
PMID:24564204
30. Huang YC, Chen TB, Hsu CC, Li SH, Wang PW, Lee BF, Kuo CY, Chiu NT. The relationship between brown adipose tissue activity and neoplastic status: an (18)F-FDG PET/CT study in the tropics. *Lipids Health Dis*. 2011; 10:238.
<https://doi.org/10.1186/1476-511X-10-238>
PMID:22182284
 31. Rousseau C, Bourbouloux E, Campion L, Fleury N, Bridji B, Chatal JF, Resche I, Campone M. Brown fat in breast cancer patients: analysis of serial (18)F-FDG PET/CT scans. *Eur J Nucl Med Mol Imaging*. 2006; 33:785–91.
<https://doi.org/10.1007/s00259-006-0066-x>
PMID:16596378
 32. Singh R, Parveen M, Basgen JM, Fazel S, Meshesha MF, Thames EC, Moore B, Martinez L, Howard CB, Vergnes L, Reue K, Pervin S. Increased Expression of Beige/Brown Adipose Markers from Host and Breast Cancer Cells Influence Xenograft Formation in Mice. *Mol Cancer Res*. 2016; 14:78–92.
<https://doi.org/10.1158/1541-7786.MCR-15-0151>
PMID:26464213
 33. Tayanloo-Beik A, Nikkhah A, Alaei S, Goodarzi P, Rezaei-Tavirani M, Mafi AR, Larijani B, Shouroki FF, Arjmand B. Brown adipose tissue and alzheimer's disease. *Metab Brain Dis*. 2023; 38:91–107.
<https://doi.org/10.1007/s11011-022-01097-z>
PMID:36322277
 34. O'Shaughnessy CT, Rothwell NJ, Shrewsbury-Gee J. Sympathetically mediated hypermetabolic response to cerebral ischemia in the rat. *Can J Physiol Pharmacol*. 1990; 68:1334–7.
<https://doi.org/10.1139/y90-201>
PMID:2078825
 35. Chao CM, Chen CL, Niu KC, Lin CH, Tang LY, Lin LS, Chang CP. Hypobaric hypoxia preconditioning protects against hypothalamic neuron apoptosis in heat-exposed rats by reversing hypothalamic overexpression of matrix metalloproteinase-9 and ischemia. *Int J Med Sci*. 2020; 17:2622–34.
<https://doi.org/10.7150/ijms.47560>
PMID:33162790
 36. Vatner DE, Oydanich M, Zhang J, Campbell SC, Vatner SF. Exercise enhancement by RGS14 disruption is mediated by brown adipose tissue. *Aging Cell*. 2023; 22:e13791.
<https://doi.org/10.1111/acer.13791>
PMID:36905127
 37. Gong B, Tang J, Jiang X, Zhang Z, Li S, Jin H, Nie L, Huang G. In situ fluorescence-photoacoustic measurement of the changes of brown adipose tissue in mice under hindlimb unloading. *J Appl Physiol* (1985). 2023; 135:251–9.
<https://doi.org/10.1152/jappphysiol.00212.2023>
PMID:37318984
 38. Tsuji T, Bussberg V, MacDonald AM, Narain NR, Kiebish MA, Tseng YH. Transplantation of Brown Adipose Tissue with the Ability of Converting Omega-6 to Omega-3 Polyunsaturated Fatty Acids Counteracts High-Fat-Induced Metabolic Abnormalities in Mice. *Int J Mol Sci*. 2022; 23:5321.
<https://doi.org/10.3390/ijms23105321>
PMID:35628137
 39. Darcy J, Tseng YH. CombATING aging-does increased brown adipose tissue activity confer longevity? *Geroscience*. 2019; 41:285–96.
<https://doi.org/10.1007/s11357-019-00076-0>
PMID:31230192
 40. Stanford KI, Middelbeek RJ, Townsend KL, An D, Nygaard EB, Hitchcox KM, Markan KR, Nakano K, Hirshman MF, Tseng YH, Goodyear LJ. Brown adipose tissue regulates glucose homeostasis and insulin sensitivity. *J Clin Invest*. 2013; 123:215–23.
<https://doi.org/10.1172/JCI62308>
PMID:23221344
 41. Peres Valgas da Silva C, Shettigar VK, Baer LA, Abay E, Madaris KL, Mehling MR, Hernandez-Saavedra D, Pinckard KM, Seculov NP, Ziolo MT, Stanford KI. Brown adipose tissue prevents glucose intolerance and cardiac remodeling in high-fat-fed mice after a mild myocardial infarction. *Int J Obes (Lond)*. 2022; 46:350–8.
<https://doi.org/10.1038/s41366-021-00999-9>
PMID:34716427
 42. Mattson MP. Perspective: Does brown fat protect against diseases of aging? *Ageing Res Rev*. 2010; 9:69–76.
<https://doi.org/10.1016/j.arr.2009.11.004>
PMID:19969105
 43. Kopelman PG. Obesity as a medical problem. *Nature*. 2000; 404:635–43.
<https://doi.org/10.1038/35007508>
PMID:10766250
 44. Han JC, Lawlor DA, Kimm SY. Childhood obesity. *Lancet*. 2010; 375:1737–48.
[https://doi.org/10.1016/S0140-6736\(10\)60171-7](https://doi.org/10.1016/S0140-6736(10)60171-7)
PMID:20451244
 45. Martin AR, Chung S, Koehler K. Is Exercise a Match for Cold Exposure? Common Molecular Framework for Adipose Tissue Browning. *Int J Sports Med*. 2020; 41:427–42.
<https://doi.org/10.1055/a-1100-7118>
PMID:32252102

46. Roubenoff R. Sarcopenia and its implications for the elderly. *Eur J Clin Nutr.* 2000 (Suppl 3); 54:S40–7.
<https://doi.org/10.1038/sj.ejcn.1601024>
PMID:11041074
47. Gonzalez-Freire M, de Cabo R, Bernier M, Sollott SJ, Fabbri E, Navas P, Ferrucci L. Reconsidering the Role of Mitochondria in Aging. *J Gerontol A Biol Sci Med Sci.* 2015; 70:1334–42.
<https://doi.org/10.1093/gerona/glv070>
PMID:25995290
48. Joseph AM, Adhihetty PJ, Buford TW, Wohlgemuth SE, Lees HA, Nguyen LM, Aranda JM, Sandesara BD, Pahor M, Manini TM, Marzetti E, Leeuwenburgh C. The impact of aging on mitochondrial function and biogenesis pathways in skeletal muscle of sedentary high- and low-functioning elderly individuals. *Aging Cell.* 2012; 11:801–9.
<https://doi.org/10.1111/j.1474-9726.2012.00844.x>
PMID:22681576
49. Short KR, Bigelow ML, Kahl J, Singh R, Coenen-Schimke J, Raghavakaimal S, Nair KS. Decline in skeletal muscle mitochondrial function with aging in humans. *Proc Natl Acad Sci U S A.* 2005; 102:5618–23.
<https://doi.org/10.1073/pnas.0501559102>
PMID:15800038
50. Coen PM, Jubrias SA, Distefano G, Amati F, Mackey DC, Glynn NW, Manini TM, Wohlgemuth SE, Leeuwenburgh C, Cummings SR, Newman AB, Ferrucci L, Toledo FG, et al. Skeletal muscle mitochondrial energetics are associated with maximal aerobic capacity and walking speed in older adults. *J Gerontol A Biol Sci Med Sci.* 2013; 68:447–55.
<https://doi.org/10.1093/gerona/gls196>
PMID:23051977
51. Conley KE, Jubrias SA, Cress ME, Esselman P. Exercise efficiency is reduced by mitochondrial uncoupling in the elderly. *Exp Physiol.* 2013; 98:768–77.
<https://doi.org/10.1113/expphysiol.2012.067314>
PMID:23085769
52. Garatachea N, Pareja-Galeano H, Sanchis-Gomar F, Santos-Lozano A, Fiuza-Luces C, Morán M, Emanuele E, Joyner MJ, Lucia A. Exercise attenuates the major hallmarks of aging. *Rejuvenation Res.* 2015; 18:57–89.
<https://doi.org/10.1089/rej.2014.1623>
PMID:25431878
53. Liu CJ, Latham NK. Progressive resistance strength training for improving physical function in older adults. *Cochrane Database Syst Rev.* 2009; 2009:CD002759.
<https://doi.org/10.1002/14651858.CD002759.pub2>
PMID:19588334
54. Serra-Rexach JA, Bustamante-Ara N, Hierro Villarán M, González Gil P, Sanz Ibáñez MJ, Blanco Sanz N, Ortega Santamaría V, Gutiérrez Sanz N, Marín Prada AB, Gallardo C, Rodríguez Romo G, Ruiz JR, Lucia A. Short-term, light-to moderate-intensity exercise training improves leg muscle strength in the oldest old: a randomized controlled trial. *J Am Geriatr Soc.* 2011; 59:594–602.
<https://doi.org/10.1111/j.1532-5415.2011.03356.x>
PMID:21453381
55. Sullivan DH, Roberson PK, Smith ES, Price JA, Bopp MM. Effects of muscle strength training and megestrol acetate on strength, muscle mass, and function in frail older people. *J Am Geriatr Soc.* 2007; 55:20–8.
<https://doi.org/10.1111/j.1532-5415.2006.01010.x>
PMID:17233681
56. Ellis T, Motl RW. Physical activity behavior change in persons with neurologic disorders: overview and examples from Parkinson disease and multiple sclerosis. *J Neurol Phys Ther.* 2013; 37:85–90.
<https://doi.org/10.1097/NPT.0b013e31829157c0>
PMID:23632452
57. Meeusen R. Exercise, nutrition and the brain. *Sports Med.* 2014 (Suppl 1); 44:S47–56.
<https://doi.org/10.1007/s40279-014-0150-5>
PMID:24791916
58. Kajimura S, Spiegelman BM, Seale P. Brown and Beige Fat: Physiological Roles beyond Heat Generation. *Cell Metab.* 2015; 22:546–59.
<https://doi.org/10.1016/j.cmet.2015.09.007>
PMID:26445512
59. Oh-ishi S, Kizaki T, Toshinai K, Haga S, Fukuda K, Nagata N, Ohno H. Swimming training improves brown-adipose-tissue activity in young and old mice. *Mech Ageing Dev.* 1996; 89:67–78.
[https://doi.org/10.1016/0047-6374\(96\)01727-7](https://doi.org/10.1016/0047-6374(96)01727-7)
PMID:8844640
60. Yoshioka K, Yoshida T, Wakabayashi Y, Nishioka H, Kondo M. Effects of exercise training on brown adipose tissue thermogenesis in ovariectomized obese rats. *Endocrinol Jpn.* 1989; 36:403–8.
<https://doi.org/10.1507/endocrj1954.36.403>
PMID:2555143
61. Slusher AL, Whitehurst M, Zoeller RF, Mock JT, Maharaj M, Huang CJ. Attenuated fibroblast growth factor 21 response to acute aerobic exercise in obese individuals. *Nutr Metab Cardiovasc Dis.* 2015; 25:839–45.
<https://doi.org/10.1016/j.numecd.2015.06.002>
PMID:26141939
62. Wu MV, Bikopoulos G, Hung S, Ceddia RB. Thermogenic capacity is antagonistically regulated in

- classical brown and white subcutaneous fat depots by high fat diet and endurance training in rats: impact on whole-body energy expenditure. *J Biol Chem*. 2014; 289:34129–40.
<https://doi.org/10.1074/jbc.M114.591008>
 PMID: [25344623](#)
63. De Matteis R, Lucertini F, Guescini M, Polidori E, Zeppa S, Stocchi V, Cinti S, Cuppini R. Exercise as a new physiological stimulus for brown adipose tissue activity. *Nutr Metab Cardiovasc Dis*. 2013; 23:582–90.
<https://doi.org/10.1016/j.numecd.2012.01.013>
 PMID: [22633794](#)
64. Tanaka R, Fuse S, Kuroiwa M, Amagasa S, Endo T, Ando A, Kime R, Kurosawa Y, Hamaoka T. Vigorous-Intensity Physical Activities Are Associated with High Brown Adipose Tissue Density in Humans. *Int J Environ Res Public Health*. 2020; 17:2796.
<https://doi.org/10.3390/ijerph17082796>
 PMID: [32325644](#)
65. Motiani P, Virtanen KA, Motiani KK, Eskelinen JJ, Middelbeek RJ, Goodyear LJ, Savolainen AM, Kempainen J, Jensen J, Din MU, Saunavaara V, Parkkola R, Löyttyniemi E, et al. Decreased insulin-stimulated brown adipose tissue glucose uptake after short-term exercise training in healthy middle-aged men. *Diabetes Obes Metab*. 2017; 19:1379–88.
<https://doi.org/10.1111/dom.12947>
 PMID: [28318098](#)
66. Singhal V, Maffazioli GD, Ackerman KE, Lee H, Elia EF, Woolley R, Kolodny G, Cypess AM, Misra M. Effect of Chronic Athletic Activity on Brown Fat in Young Women. *PLoS One*. 2016; 11:e0156353.
<https://doi.org/10.1371/journal.pone.0156353>
 PMID: [27243823](#)
67. Vosselman MJ, Hoeks J, Brans B, Pallubinsky H, Nascimento EB, van der Lans AA, Broeders EP, Mottaghy FM, Schrauwen P, van Marken Lichtenbelt WD. Low brown adipose tissue activity in endurance-trained compared with lean sedentary men. *Int J Obes (Lond)*. 2015; 39:1696–702.
<https://doi.org/10.1038/ijo.2015.130>
 PMID: [26189600](#)
68. Liu X, Zheng Z, Zhu X, Meng M, Li L, Shen Y, Chi Q, Wang D, Zhang Z, Li C, Li Y, Xue Y, Speakman JR, Jin W. Brown adipose tissue transplantation improves whole-body energy metabolism. *Cell Res*. 2013; 23:851–4.
<https://doi.org/10.1038/cr.2013.64>
 PMID: [23649313](#)